

## Synthesis of perylene-bridged bis(dibenzo-24-crown-8) and its assembly behavior with a fullerene-based secondary dibenzylammonium salt

YAO HaiQing, ZHANG HengYi, HAN Min, DING ZhiJun, ZHANG ZhiJun & LIU Yu\*

*State Key Laboratory of Elemento-Organic Chemistry; Department of Chemistry, Nankai University, Tianjin 300071, China*

Received April 29, 2010; accepted May 11, 2010

This paper reports the synthesis of a perylene-bridged bis(crown ether) through the condensation reaction between primary amine-armed dibenzo-24-crown-8 and 3,4,9,10-perylenetetracarboxylic dianhydride. Pseudorotaxane with a perylene moiety at the wheel and a fullerene unit in the middle of the axle was prepared, which was evidenced by  $^1\text{H}$  NMR and fluorescence experiments. Subsequent investigations show that the formation and disassociation of the pseudorotaxane can be controlled by the alternating addition of  $\text{KPF}_6$  and 18-crown-6, following the change of the fluorescence intensity of perylene.

**crown ether, perylene, fullerene, pseudorotaxane**

### 1 Introduction

Since 1950, perylene bisimide derivatives have found high grade industrial applications [1] as pigments due to their favorable combination of insolubility and migrational stability, light- and weather-fastness, thermal stability and chemical inertness. Recently, perylene derivatives as n-type semiconductors are also found to be promising compounds for applications in optical [2] and electronic devices [3], such as electrophotography [4], organic field effect transistors [5], photovoltaic devices [6], and organic solar cells [7]. The intrinsic insolubility of perylene bisimides is the main barrier for synthesis, purification and application. To prepare soluble perylene bisimide dyes, Langhals [8] and Seybold *et al.* [9] provided two successful strategies. The first way is to introduce solubilizing substituents at the imide nitrogen, and the other way is to introduce substituents at the carbocyclic scaffold in the so-called bay-area.

Fullerene  $\text{C}_{60}$  possesses unique and attractive electronic

and optical properties owing to the ability of remarkable acceleration of photoinduced charge separation and deceleration of charge recombination in donor-fullerene systems [10]. In order to construct fullerene-containing supramolecular assemblies and advanced materials, fullerene has been incorporated into various systems such as porphyrins [11], ferrocene [12], and hydroquinone [13]. The self-assembly of fullerene-based molecules has aroused much interest in the construction of more complex two- and three-dimensional systems [14].

Dibenzo-24-crown-8 (DB24C8) can form 1:1 complexes with the dialkylammonium cations. This unique binding model has been utilized to construct various supramolecular assemblies containing DB24C8 and dialkylammonium salts [15]. Recently, we reported a reversible luminescent lanthanide switch based on this binding model [16] and a bistable [3] rotaxane with intramolecular charge-transfer behavior [17]. Herein, we synthesized a new perylene-bridged bis-(DB24C8) host molecule (**1**), and constructed a pseudorotaxane system including **1** and a fullerene-based secondary dibenzylammonium cation (**2**) (Figure 1). The reversible fluorescence process of the pseudorotaxane can be modulated

\*Corresponding author (email: yuliu@nankai.edu.cn)




Figure 1 Structure of host **1** and guest **2**.

by controlling the binding and release of the guest molecule **2**.

## 2 Experimental

### 2.1 Materials and apparatus

3,4,9,10-Perylene dianhydride (Sigma-Aldrich company), 4-carboxylic-aciddibenzo-24-crown-8 (**3**) [18], compounds **6** [19], **7** [19] and the fullerene-based secondary dibenzylammonium salt (**2**) [20] were prepared according to the previously reported procedure. Chloroform and dichloromethane were redistilled and kept with 4 Å molecular sieve. Acetonitrile is refluxed for 10 h in the presence of  $K_2CO_3$  and  $KMnO_4$ , stirred with a proper amount of  $P_2O_5$  overnight, and then distilled at 81–82 °C. All the other chemicals were commercially available.

$^1H$  NMR spectra were recorded on a Varian Mercury VX400 spectrometer. Fluorescence spectra were measured with JASCO FP-750 spectrometer. ESI-MS was performed on Thermofinnigan LCQ Advantage LC-MS. MALDI-TOF was performed on Bruker Autoflex III.

### 2.2 Synthesis of host **1**

*tert*-Butyl 2-(6,7,9,10,12,13,20,21,23,24,26,27-dodecahydrodibenzo[*b,n*][1,4,7,10,13,16,19,22]octaoxacyclotetracosine-2-carboxamido)ethylcarbamate (**4**)

4-Carboxylic-aciddibenzo-24-crown-8 (**3**) (2.0 g, 4.1 mmol), *tert*-butyl-2-aminoethyl carbamate (0.7 g, 4.6 mmol), *N,N'*-dicyclohexylcarbodiimide (1.4 g, 6.8 mmol) and a little of *N,N*-dimethylamino pyridine were dissolved in dry  $CHCl_3$  and the mixture was stirred in an ice bath for 12 h. The reaction mixture was filtered. The filtrate was washed with water twice, and dried over anhydrous sodium sulfate. Then the solvent was removed at reduced pressure to give a white residue. The residue was purified by silica-gel column chromatography using  $CHCl_3/CH_3OH$  (20/1) (*v/v*) as the eluent to give product **4** (1.4 g, 50%);  $^1H$  NMR (400 MHz,  $CDCl_3$ , 298 K)  $\delta$  1.42 (s, 9H), 3.40 (t, 2H), 3.52 (t, 2H),

3.83 (m, 8H), 3.92 (m, 8H), 4.18 (m, 8H), 6.88 (d, 1H), 6.88 (m, 4H), 7.33 (dd, 1H), 7.42 (d, 1H),  $m/z$  657  $[M+Na]^+$ .

*N*-(2-Aminoethyl)-6,7,9,10,12,13,20,21,23,24,26,27-dodecahydrodibenzobenzene[1,4,7,10,13,16,19,22]octaoxacyclo-tetracosine-2-carboxamide (**5**)

Compound **4** was dissolved in 10 mL  $CHCl_3$ , and the mixture was stirred at room temperature. 4 mL of  $CF_3COOH$  was added dropwise. The stirring was continued overnight and followed by evaporation of the solvent. The residue was washed with hexane for three times. The crude product was dissolved in  $CH_3OH$  and the pH of the solution was adjusted to 8 by NaOH solution. The crude product was purified by silica-gel column chromatography using  $CHCl_3/CH_3OH$  (5/1) (*v/v*) as the eluent to give product **5** (0.5 g, 90%);  $^1H$  NMR (400 MHz,  $CDCl_3$ , 298 K)  $\delta$  2.95 (t, 2H), 3.48 (t, 2H), 3.82 (m, 8H), 3.92 (m, 8H), 4.17 (m, 8H), 6.83 (d, 1H), 6.88 (m, 4H), 7.30 (dd, 1H), 7.41 (d, 1H),  $m/z$  557  $[M+Na]^+$ .

### Compound **1**

Compounds **7** (300 mg, 0.3 mmol) and **5** (500 mg, 0.9 mmol) were dissolved in 6 mL isopropanol, and then 0.1 mL triethylamine was added. The mixture was refluxed under  $N_2$  atmosphere for two days. The mixture was cooled to room temperature and followed by evaporation of the solvent. The crude product was purified by silica-gel column chromatography using  $CHCl_3/CH_3OH$  (20/1) (*v/v*) as the eluent to give red powder **1** (300 mg, 50%);  $^1H$  NMR (400 MHz,  $CDCl_3$ , 298 K)  $\delta$  1.28 (s, 36H), 3.67 (m, 12H), 3.72 (m, 12H), 3.82 (m, 12H), 4.02 (m, 12H), 4.28 (m, 4H), 6.78 (d, 8H) 6.85 (m, 8H), 7.18 (m, 4H), 7.25 (d, 8H), 7.50 (s, 4H), 7.99 (s, 4H)  $^{13}C$  NMR (400 MHz,  $CDCl_3$ , 298 K)  $\delta$  167.11, 164.16, 155.92, 152.89, 151.35, 148.85, 148.55, 147.34, 132.85, 127.17, 126.68, 122.09, 121.41, 120.90, 120.30, 119.62, 119.53, 119.39, 119.17, 114.01, 112.59, 71.35, 71.22, 71.16, 69.90, 69.68, 69.32, 69.23, 69.12, 34.38, 31.46. MS (MALDI-TOF)  $m/z$ : Anal. calcd  $C_{118}H_{128}O_{26}N_4$ :  $[M+Na]^+$ : 2040.8743, found  $[M+Na]^+$ : 2040.8766.

### 3 Results and discussion

#### 3.1 Synthesis

The synthetic pathway of **1** is shown in Figure 2. Compound **5** was obtained by DCC condensation of 4-carboxylic-ciddibenzo-24-crown-8 (**3**) with tert-butyl-2-aminoethyl carbamate, followed by CF<sub>3</sub>COOH reduction of the Boc-protected group. 3,4,9,10-Perylene tetracarboxylic dianhydride is insoluble in organic solvent. To improve its solubility, its bay-area was substituted by four chlorine atoms by reacting with HSO<sub>3</sub>Cl, the propyl substituents were introduced through reaction with propylamine. Compound **6** was obtained by introducing four *tert*-butyl phenols at the bay-area. Subsequent hydrolyzation of **6** with a base produced compound **7**. Finally, the target compound **1** was obtained by the reaction of **5** with **7** in isopropanol using triethylamine as the base. Owing to those soluble substitutions, compound **1** possesses very good solubility in organic solvent.

#### 3.2 <sup>1</sup>H NMR spectra of the complex of **1** with **2**


To obtain in-depth mechanistic insight into the complexation between host **1** and guest **2**, we performed <sup>1</sup>H NMR experiments. Figure 3 shows the signals of some protons moved apparently upon complexation. The Ha proton in **2** shifted downfield from 3.90 to 4.62 ppm. This may be attributed to the hydrogen bond interaction between Ha and oxygen atoms of the crown ether. The resonances of the Hc proton in **2** split into two equal parts. A half of them shifted upfield, while the other remained unchanged. This means that a half of **2** was located in the ring of the crown ether, while the other did not bind with **1**. In addition, the signals of protons in crown ethers (3.4–4.5 ppm) and perylene (8.06 ppm) split apparently, indicating that the distance between C<sub>60</sub> and the perylene unit became close after complexation. These spectral changes validate the formation of the pseudorotaxane [**1+2**].

#### 3.3 Fluorescence spectra of the interaction of **1** with **2**


Benefiting from the fascinating fluorescence property of perylene, the host **1** displays satisfactory luminescent emission in a CHCl<sub>3</sub>:CH<sub>3</sub>CN = 2:1 (v/v) solution (Figure 4). When excited at 510 nm, **1** showed two emission peaks at 620 nm (0→0) and a shoulder peak at 675 nm (0→1) [21]. When 22 equiv of the guest **2** was added to the solution, the fluorescence emission at 620 nm of **1** was significantly quenched to 75% of its original intensity. Instead of **2** with a compound bearing the C<sub>60</sub> moiety but without secondary dibenzyl-ammonium unit, the luminescence of **1** slightly decreased, which is attributed to the absorption of C<sub>60</sub> existing at 450–750 nm. In addition, dibenzylammonium hexafluorophosphate does not influence the luminescence of **1**. These combined observations suggest that both the C<sub>60</sub> unit and the dibenzylammonium cation are two prerequisites to quench the fluorescence emission of **1**. That indicates the interaction of **1** with **2** leads to the formation of the pseudorotaxane [**1+2**], so a PET process from perylene of **1** to the C<sub>60</sub> moiety of **2** occurs due to the adjacency of C<sub>60</sub> with perylene.

#### 3.4 Reversible fluorescence switch


The binding behavior of 24C8 and the dialkylammonium cation can be switched reversibly by various external stimuli, such as pH changes and competition binding. The association constants of DB24C8 with K<sup>+</sup> and dialkylammonium cations are 7.6 × 10<sup>3</sup> and 1.2 × 10<sup>3</sup> M<sup>-1</sup>, respectively, while that of 18-crown-6 (18C6) with K<sup>+</sup> is 1.3 × 10<sup>6</sup> M<sup>-1</sup> [21]. From those data, we might deduce reasonably that the dialkylammonium cation in DB24C8 could be replaced by K<sup>+</sup> upon the addition of KPF<sub>6</sub> to the solution of the pseudorotaxane [**1+2**], and then the supramolecular assembly could be reproduced upon the addition of 18-crown-6. To evidence this assumption, we performed the fluorescent experiments of the pseudorotaxane [**1+2**] in the absence and




**Figure 2** Synthetic routes to host **1**. (a) DCC, DMAP, BocNHCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CHCl<sub>3</sub>, 12 h, 50%. (b) TFA, 12 h, 90%. (c) (1) HSO<sub>3</sub>Cl, I<sub>2</sub>, 20 h, 40%; (2) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, H<sub>2</sub>O/CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH, 8 h, 99%. (3) ArOH, KCO<sub>3</sub>, NMP, 8 h, 90%. (d) KOH/H<sub>2</sub>O, CH<sub>3</sub>CHOHCH<sub>3</sub>, 12 h, 78%. (e) N(Et)<sub>3</sub>, CH<sub>3</sub>CHOHCH<sub>3</sub>, 2 d, 50%.



**Figure 3** Partial  $^1\text{H}$  NMR spectra of (400 MHz,  $\text{CD}_3\text{CN}/\text{CDCl}_3=1/2$ , 298 K,  $[\mathbf{1}]_0=3$  mM) (a)  $\mathbf{1}$ , (b)  $\mathbf{1}$  and 2.0 equiv of  $\mathbf{2}$ , and (c)  $\mathbf{2}$  (Hcu: peak of the uncomplex; Hcc: peak of the complex; \*: peak of  $\text{CDCl}_3$ ).



**Figure 4** The fluorescence spectra of compound  $\mathbf{1}$  with increasing concentration of  $\mathbf{2}$  ( $\text{CHCl}_3:\text{CH}_3\text{CN}=2:1$ , 295 K);  $[\mathbf{1}]=1\times 10^{-5}$  M and  $[\mathbf{2}]=1\times 10^{-5}$  to  $2.2\times 10^{-4}$  M.  $\lambda_{\text{exc}}=510$  nm.



**Figure 5** Emission spectral changes observed for  $\mathbf{1}$ ,  $[\mathbf{1}+\mathbf{2}]$ ,  $[\mathbf{1}+\mathbf{2}]+\text{K}^+$ , and  $[\mathbf{1}+\mathbf{2}]+\text{K}^++18\text{-crown-6}$ . The emission changes of  $\mathbf{1}$  at 620 nm (inset) in the presence ( $\mathbf{1}$ ) of  $\mathbf{2}$  and added  $\text{KPF}_6$  (4, 6, 8, 10) and 18-crown-6 (3, 5, 7, 9) in  $\text{CH}_3\text{CN}/\text{CHCl}_3$  solution (1:2).

presence of  $\text{KPF}_6$  and 18-crown-6.

As shown in Figure 5, when  $\text{KPF}_6$  was added to the solution of the pseudorotaxane  $[\mathbf{1}+\mathbf{2}]$ , the quenched perylene

emission was restored (91%). This observation suggests that the dialkylammonium cation in DB24C8 was replaced by  $\text{K}^+$  and the PET process from the perylene moiety to  $\text{C}_{60}$  was suppressed. To check the reversibility of this process, 18-crown-6 was added to the solution. The luminescence quenching was reproduced again as a proof of regeneration of the pseudorotaxane supramolecular assembly. The process can be repeated several times. These experiments demonstrate that the fluorescence of the pseudorotaxane  $[\mathbf{1}+\mathbf{2}]$  can be switched in a reversible manner by external chemical stimuli.

## 4 Conclusions

In conclusion, a perylene-bridged bis(dibenzo-24-crown-8)  $\mathbf{1}$  and a guest molecule  $\mathbf{2}$  containing  $\text{C}_{60}$  unit have been synthesized. The binding behavior and the photophysical behavior of the resultant pseudorotaxane between  $\mathbf{1}$  and  $\mathbf{2}$  have been investigated through  $^1\text{H}$  NMR and fluorescence spectra. The results obtained indicate that the fluorescence of  $\mathbf{1}$  was quenched significantly in the presence of  $\mathbf{2}$ , which is attributed to a PET from perylene of  $\mathbf{1}$  to the  $\text{C}_{60}$  moiety of  $\mathbf{2}$  in this pseudorotaxane system. The reversible fluorescence process can be modulated by controlling the binding and release of the guest molecules by adding  $\text{K}^+$  or 18C6, respectively. It is an effective strategy to control the fluorescent behavior of perylene through external chemical stimuli.

This work was supported by the National Basic Research Program of China (973 Program) (2006CB932900) and the National Natural Science Foundation of China (20932004 and 20772063).

- 1 Würthner F. Perylene bisimide dyes as versatile building blocks for functional supramolecular architectures. *Chem Commun*, 2004, 14: 1564–1579
- 2 Yan LW, Yang L, Lan JB, You JS. A new perylene diimide-based colorimetric and fluorescent sensor for selective detection of  $\text{Cu}^{2+}$  cation. *Sci China Ser B-Chem*, 2009, 52(4): 518–522
- 3 Struijk CW, Sieval AB, Dakhorst JE, Dijk MV, Kimkes P, Koehorst RB, Donker H, Schaafsma TJ, Picken SJ, Craats AM, Warman JM, Zuilhof H, Sudhölter EJ. Liquid crystalline perylene diimides: Architecture and charge carrier mobilities. *J Am Chem Soc*, 2000, 122: 11057–11066
- 4 Law KY. Organic photoconductive materials: Recent trends and developments. *Chem Rev*, 1993, 93: 449–486
- 5 (a) Jones BA, Ahrens MJ, Yoon MH, Facchetti A, Marks TJ, Wasielewski MR. High-mobility air-stable *n*-type semiconductors with processing versatility: Dicyanoperylene-3,4,9,10-bis(dicarboximides). *Angew Chem Int Ed*, 2004, 43: 6363–6366; (b) Briseno AL, Mannsfeld SC, Reese C, Hancock JM, Xiong YJ, Jenekhe SA, Bao ZN, Xia YN. Perylenediimide nanowires and their use in fabricating field-effect transistor and complementary inverters. *Nano Lett*, 2007, 7: 2847–2853
- 6 (a) Hua JL, Meng FS, Ding F, Li FY, Tian H. Novel soluble and thermally-stable fullerene dyad containing perylene. *J Mater Chem*, 2004, 14: 1849–1853; (b) Liu Y, Wang N, Li YJ, Liu HB, Li YL, Xiao JC, Xu XH, Huang CS, Cui S, Zhu DB. A new class of conju-

- gated polyacetylenes having perylene bisimide units and pendant fullerene or porphyrin groups. *Macromolecules*, 2005, 38: 4880–4887
- 7 (a) Schmidt-Mende L, Fechtenkötter A, Mullen K, Moons E, Friend RH, MacKenzie JD. Self-organized discotic liquid crystals for high-efficiency organic photovoltaics. *Science*, 2001, 293: 1119–1122; (b) Li JL, Kastler M, Pisula W, Robertson JW, Wasserfallen D, Grimsdale AC, Wu JS, Müllen K. Organic bulk-heterojunction photovoltaics based on alkyl substituted discotics. *Adv Funct Mater*, 2007, 17: 2528–2533; (c) Pandey AK, Nunzi JM. Upconversion injection in rubrene/perylene-diimide-heterostructure electroluminescent diodes. *Appl Phys Lett*, 2007, 90: 263508, 1–3
- 8 Langhals H. Cyclic carboxylic imide structures as structure elements of high stability. Novel developments in perylene dye chemistry. *Heterocycles*, 1995, 40: 477–500
- 9 (a) Seybold G, Wagenblast G. New perylene and violanthrone dye-stuffs for fluorescent collectors. *Dyes Pigm*, 1989, 11: 303–317; (b) Seybold G, Stange A (BASF AG), Patent, DE 3545004, 1987-12-19. (*Chem Abstr*, 1988, 108: 77134c)
- 10 (a) Reed CA, Bolskar RD. Discrete fulleride anions and fullerenium cations. *Chem Rev*, 2000, 100: 1075–1120; (b) Gust D, Moore TA, Moore AL. Mimicking photosynthetic solar energy transduction. *Acc Chem Res*, 2001, 34: 40–48
- 11 (a) Shirakawa M, Fujita N, Shinkai S. [60]Fullerene-motivated organogel formation in a porphyrin derivative bearing programmed hydrogen-bonding sites. *J Am Chem Soc*, 2003, 125: 9902–9903; (b) Yamaguchi T, Ishii N, Tashiro K, Aida T. Supramolecular peapods composed of a metalloporphyrin nanotube and fullerenes. *J Am Chem Soc*, 2003, 125: 13934–13935
- 12 Crane JD, Hitchcock PB, Kroto HW, Taylor R, Walton DR. Preparation and characterisation of  $C_{60}$ (ferrocene)<sub>2</sub>. *J Chem Soc, Chem Commun*, 1992, 24: 1764–1765
- 13 Diederich F, Gómez-López M. Supramolecular fullerene chemistry. *Chem Soc Rev*, 1999, 28:263–277, and references therein
- 14 (a) Kimura M, Saito Y, Ohta K, Hanabusa K, Shirai H, Kobayashi N. Self-organization of supramolecular complex composed of rigid dendritic porphyrin and fullerene. *J Am Chem Soc*, 2002, 124: 5274–5275; (b) Hasobe T, Kashiwagi Y, Absalom MA, Sly J, Hosomizu K, Crossley MJ, Imahori H, Kamat PV, Fukuzumi S. Supramolecular photovoltaic cells using porphyrin dendrimers and fullerenes. *Adv Mater*, 2004, 16: 975–979; (c) Sun D, Tham FS, Reed CA, Chaker L, Boyd PD. Supramolecular fullerene-porphyrin chemistry. Fullerene complexation by metalated “Jaws Porphyrin” hosts. *J Am Chem Soc*, 2002, 124: 6604–6612
- 15 (a) Zhu XZ, Chen CF. A highly efficient approach to [4]pseudocatenanes by threefold metathesis reactions of a triptycene-based tris[2]pseudorotaxane. *J Am Chem Soc*, 2005, 127: 13158–13159; (b) Wang F, Han CY, He CL, Zhou QZ, Zhang JQ, Wang C, Li N, Huang FH. Self-sorting organization of two heteroditopic monomers to supramolecular alternating copolymers. *J Am Chem Soc*, 2008, 130: 11254–11255; (c) Badjić JD, Nelson A, Cantrill SJ, Turnbull B, Stoddart JF. Multivalency and cooperativity in supramolecular chemistry. *Acc Chem Res*, 2005, 38: 723–732; (d) Zhai CX, Huang FH. Recent research progresses of pseudorotaxanes and rotaxanes (in Chinese). *Sci China Ser B-Chem*, 2009, 39(4): 315–328
- 16 Han M, Zhang HY, Yang LX, Jiang Q, Liu Y. A reversible luminescent lanthanide switch based on a dibenzo[24]-crown-8-dipicolinic acid conjugate. *Org Lett*, 2008, 10: 5557–5560
- 17 Jiang Q, Zhang HY, Han M, Ding ZJ, Liu Y. pH-Controlled intramolecular charge-transfer behavior in bistable [3]rotaxane. *Org Lett*, 2010, 12: 1728–1731
- 18 Feng DJ, Li XQ, Wang XZ, Jiang XK, Li ZT. Highly stable pseudo[2]rotaxanes co-driven by crown ether-ammonium and donor-acceptor interactions. *Tetrahedron*, 2004, 60: 6137–6144
- 19 (a) Würthner F, Thalacker C, Sautter A. Substituted diazadibenzoperylenes: new functional building blocks for supramolecular chemistry. *Angew Chem Int Ed*, 2000, 39: 1243–1245; (b) Gvishi R, Reisfeld R, Burshtein Z. Spectroscopy and laser action of the “red perylimide dye” in various solvents. *Chem Phys Lett*, 1993, 213: 338–344
- 20 Diederich F, Echegoyen L, Gómez-López M, Kessinger R, Stoddart JF. The self-assembly of fullerene-containing [2]pseudorotaxanes: Formation of a supramolecular  $C_{60}$  dimer. *J Chem Soc, Perkin Trans. 2*, 1999, 8: 1577–1586
- 21 Jérôme Baffreau, Lucie Ordonneau, Stéphanie Leroy-Lhez, and Piérick Hudhomme. Synthesis of perylene-3,4-mono(dicarboximide)-fullerene  $C_{60}$  dyads as new light-harvesting systems. *J Org Chem*, 2008, 73: 6142–6147
- 22 Frensdorff HK. Stability constants of cyclic polyether complexes with univalent cations. *J Am Chem Soc*, 1971, 93: 600–606