

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: W. Zhou, W. Lin, Y. Chen and Y. Liu, *Chem. Sci.*, 2022, DOI: 10.1039/D2SC01770A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Supramolecular Assembly Confined Purely Organic Room Temperature Phosphorescence and Its Biological Imaging

Wei-Lei Zhou^{a,b}, Wenjing Lin^a, Yong Chen^a and Yu Liu^{a*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Purely organic room temperature phosphorescence, especially in aqueous solution, is attracting increasing attention owing to the large Stokes shift, long lifetime, low preparation cost, low toxicity, good processing performance advantages, and broad application value. This review mainly focuses on macrocyclic (cyclodextrin and cucurbituril) hosts, nanoassembly, and macromolecule (polyether) confined-driven RTP. As an optical probe, the assembly and the two-staged assembly strategy can realize the confined purely organic RTP, and achieve energy transfer and light-harvesting from fluorescence to delayed fluorescence or phosphorescence. This supramolecular assembly is widely applied for luminescent material, cell imaging, and other fields because it effectively avoids oxygen quenching. In addition, the near-infrared excitation, near-infrared emission, and in situ imaging of purely organic room temperature phosphorescence in assembled confinement materials are also prospected.

Introduction

Complex biological processes need to be monitored with optical probes for imaging and sensing that provide information on probe identity, spatial location, and concentration based on the wavelength and intensity of the signal.¹ Therefore, optical imaging plays an indispensable role in the biomedical field.² However, endogenous background fluorescence or spontaneous fluorescence from cellular organelles has been always a problem, resulting in reduced imaging sensitivity and specificity.³ Given the continuous progress and development of science and technology, the sensitivity, specificity, and achievable spatial and temporal resolution of optical imaging technology have been continuously improved.^{1a, 4} The development of chemiluminescence or Cerenkov probes eliminates the problem of auto-fluorescence because they do not require light excitation, but some problems are still inevitable.^{2i, 5} For example, chemiluminescence usually requires specific enzymes and substrates, which complicates imaging process somewhat, and Cerenkov luminescence requires radioisotope radiation to be produced, which is also associated with biological safety risks.⁶ Therefore, it is highly desirable to design new luminescence probes that can overcome the abovementioned drawbacks in biological imaging applications.

Room-temperature phosphorescence (RTP) materials with longer lifetime, larger Stokes shift, and the involvement of triplet states have garnered attention in the fields of biosensing, bioimaging, and imaging-guided surgery or therapy for providing high signal-to-noise

ratio images, because they have persistent luminescence even after removing the light source, allowing in vivo imaging without real-time external excitation.⁷ However, the RTP effect is common in inorganic compounds containing rare-earth metals or heavy metals complexes and metal–organic compounds, and the high cost and the potential cytotoxicity of metals limits their biological applications.^{7n, 8-9} As avoiding the introduction of metals, purely organic RTP materials are good candidates for RTP materials. The high bonding electrons in purely organic materials limit the material ability to decay from the triplet state through the radiation relaxation pathway, and are prone to non-radiative relaxation through heat and collision, and energy transfer from the triplet state to the triplet state of oxygen.^{7f, 7h, 7n, 7p, 8d, 10}

Scheme 1 shows the formulas for the quantum yield of phosphorescence (1), the quantum yield of intersystem crossing (2), and the lifetime of the phosphorescence (3).^{7h, 7i, 7n, 7p, 7r} According to Equations 1 and 2, to obtain high phosphorescence quantum yield, it is necessary to increase the intersystem crossing rate (k_{isc}), the phosphorescence decay rate (k_r^{Phos}), and at the same time, reduce the rate of nonradiative transitions (k_{nr}^{Phos}) and suppress the influence of quenching groups. On the one hand, k_{isc} and k_r^{Phos} can be enhanced by adjusting the molecular orbital structure by introducing abundant lone-pair electrons (including O, N, P, S, etc.), introducing heavy atoms (Cl, Br, I, S, Se, etc.), narrowing the energy gap between singlet and the triplet state (use donor–acceptor-type molecules for the charge transfer state), and increasing the degree of molecular aggregation. On the other hand, k_{nr}^{Phos} and quenching groups can be controlled by improving the rigidity of the phosphorescent structure with the introduction of numerous intermolecular interactions (hydrogen bonding, halogen bonding, ionic bonding, CH– π interaction, and π – π interaction, etc.). According to Equation 3, the long-lived RTP needs to minimize k_{nr}^{Phos} and k_r^{Phos} and avoid quencher interference. In addition, the


^a College of Chemistry, State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, P. R. China

^b College of Chemistry and Material Science, Inner Mongolia Key Laboratory of Chemistry for Nature Products and Synthesis for Functional Molecules, Innovation Team of Optical Functional Molecular Devices, Inner Mongolia Minzu University, Tongliao 028000, P. R. China

Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x



introduction of long-range charge diffusion before the emission of triplet excitons is also a feasible way to improve the lifetime of RTP. Based on Equations 1 and 3, an increase in τ_{Phos} implies a decrease in Φ_{Phos} based on the change in k_r^{Phos} . However, owing to the quenching from the collision of the oxygen dissolved in water as well as the free molecular motions for nonradiative decay processes, the most efficient purely organic RTP emissions are common in the solid-state, which make them impractical for use in biological applications. Thus, to obtain purely organic RTP in water, three important aspects should be considered: (1) the enhancement of the heavy atom effect to promote intersystem crossing (ISC); (2) a rigid environment to suppress non-radiative transitions in water; (3) avoiding the influence of quenchers such as O_2 .







Scheme 1. Schematic illustration of the Jablonski diagram of the emissive processes of phosphorescence (Φ_{Phos} : the quantum yield of phosphorescence; Φ_{isc} : the quantum yield of intersystem crossing; τ_{Phos} : the lifetime of the phosphorescence; $k_r^{Fluo.}$: the fluorescent decay rate constant; $k_{nr}^{Fluo.}$: the fluorescent nonradiative rate constant; k_{isc} : the intersystem crossing rate constant from S_1 to T_1 ; $k_r^{Phos.}$: the phosphorescent decay rate constant; $k_{nr}^{Phos.}$: the phosphorescent nonradiative rate constant).

Unlike traditional covalent interactions, the supramolecular chemistry approach mainly focuses on weak noncovalent interactions, such as hydrogen bonds, π - π stacking, hydrophobic forces, van der Waals forces, and electrostatic interactions; not only do they make the single functional molecule multi-functionalized in a simple way, but they also bring new properties by adjusting molecular conformation, vibration, energy gap, or electronic distribution.^{7f, 7h, 7i, 7m, 7n, 10b, 11} More importantly, this approach can provide an effective method for the construction of water-phase purely organic RTP. In this mini-review, we briefly introduce the research progress into supramolecular strategy for the construction of a purely organic aqueous RTP system mediated by macrocyclic hosts (cyclodextrin and cucurbituril), macromolecule (polyether) and nanoassembly confinement, as well as its application in cell imaging. In addition, we provide the key challenges and perspectives for further advances in purely organic aqueous phosphorescence. We hope that this review will bring insight and inspiration to readers and allow the future development of phosphorescence and its biological applications in water. Table 1 presents a list of the chemical structures of the phosphorescent molecules involved in this paper.

Table 1. Chemical structures of phosphorescent molecules

DOI: 10.1039/D2SC01770A

Structure	Citation	Strategy
 derivative	Figs. 1–3, 15	Macrocyclic Confinement
 derivative	Figs. 4, 6, 19	Macrocyclic and Nanoassembly Confinement
 derivative	Figs. 5, 7	Macrocyclic Confinement
 derivative	Figs. 8–14	Macrocyclic Confinement
 derivative	Figs. 16, 21	Macrocyclic and Nanoassembly Confinement
 derivative	Figs. 17, 18	Nanoassembly Confinement
 derivative	Figs. 25, 28	Macromolecular Confinement
 derivative	Figs. 26, 27	Macromolecular Confinement
 derivative	Figs. 18, 23, 24	Nanoassembly and Macromolecular Confinement
 derivative	Fig. 20	Nanoassembly Confinement

Macrocyclic Confinement



The two main water-soluble biocompatible macrocyclic host compounds, cyclodextrin (CD) and cucurbituril (CB), can provide a hydrophobic cavity for the guest molecule in aqueous solution, which could protect the luminescent guest from the influence of free oxygen, as well as restrict molecular motion for reducing nonradiative decay via host-guest interaction. CDs are a type of cyclic oligosaccharides constructed by D-glucose units through α -1,4-glucose bonds and can form stable host-guest complexes with various neutral or charged phosphorescent molecules.^{7h, 7m, 11d, 12} As early as 1982, Turro et al. achieved phosphorescent emission in nitrogen-purged aqueous solutions based on the inclusion of β -CD with 1-bromonaphthalene and 1-chloronaphthalene.¹³ Furthermore, the addition of a third component with heavy atoms to the phosphorescent inclusion complex of CD with nitrogen heterocycles or bridged biphenyl compounds could also enhance the phosphorescence effect in water.¹⁴ In another case, RTP was enhanced via the formation of the ternary complex 1-bromonaphthalene/glucosyl-modified- β -CD/alcohol, in which the RTP of 1-BrNp is effectively protected from oxygen quenching.¹⁵ When 1-BrNp is modified on the secondary OH groups of β -CD, the RTP at 530 nm is achieved in aqueous solutions through the self-inclusion of 1-BrNp into the hydrophobic cavity of β -CD to shield the oxygen.¹⁶ Meanwhile, halide alkanes have an external heavy atom enhancement effect on CD-RTP in water in the order Br > Cl >> I.¹⁷ This is because of the heavy atom effect for enhancing the spin-orbit (S-O) coupling interaction, which can further increase the ISC from the singlet excited state to the triplet state. However, if the heavy atom is too strong, k_{nr}^{Phos} may also increase greatly, and result in a significant reduction in RTP. Among them, dibromoalkanes are the best of all bromoalkanes, and *t*-butyl chloride can induce RTP of polycyclic aromatic compounds more effectively. Adding a third part of hexahydropyridine (HHP), cyclohexane (CH), or 1-ethylpiperidine (EP) to 1-BrNp/ β -CD aqueous solution could improve the RTP of 1-BrNp/ β -CD; in particular, the addition of EP exhibited the strongest RTP due to the large rigidity.¹⁸ Furthermore, the photocontrolled phosphorescence of CD-RTP in water can be realized by the assembly and disassembly of cyclodextrin and azobenzene. For example, a photocontrolled reversible RTP pseudorotaxane in aqueous solution was constructed via a β -CD/ α -BrNp/sodium 2-methoxy-5-((4-nitrophenyl)diazanyl)benzoate (DAYR) ternary system (Fig. 1).¹⁹ In this system, trans-DAYR could form a pseudorotaxane with β -CD. When the trans-DAYR converted the cis-DAYR via the photoirradiation, the DAYR/ β -CD pseudorotaxane was disassembled, and the inclusion complex of α -BrNp/ β -CD was formed, achieving RTP, because the order of bonding constants was β -CD/trans-DAYR > β -CD/ α -BrNp > β -CD/cis-DAYR. In the binary RTP system between the azo modified β -CD and α -BrNp, the RTP of β -CD/ α -BrNp could be reversibly controlled via the photoisomerization of the azo group in and out of the cyclodextrin cavity (Fig. 2).²⁰ Moreover, this regulation of phosphorescence can also be achieved in polymers. Three polymers, i.e., poly-BrNp, poly- β -CD, and poly-azo were prepared by the copolymerization of acrylamide and monomer of α -BrNp, β -CD or azobenzene respectively (Fig. 3).²¹ The weak

RTP emission was observed as the addition of β -CD into the polymer poly-BrNp aqueous solution. However, the stronger RTP emission was achieved while the poly- β -CD aqueous solution was added into the poly-BrNp aqueous solution owing to the formation of hydrogel through the host-guest interaction for protecting BrNp moiety to some extent. When aqueous solutions of three components poly-BrNp, poly- β -CD and poly-Azo were mixed to form the smart hydrogel, the photo-stimulated RTP was realized through the trans-cis photoisomerization of azobenzene units for the threading and de-threading into the cavity of β -CD. Another polycyclic aromatic hydrocarbon 4-bromo-1, 8-naphthalic anhydride (BrNpA) was used to investigate the RTP in water (Fig. 4),²² and a photo-controlled RTP supramolecular assembly was constructed in aqueous solution through the ternary poly-BrNpA/ γ -CD/poly-Azo system, where the phosphorescence excitation and emission wavelengths presented large red shifts compared with previously reported systems.




Fig. 1 Schematic illustration of photo-controlled reversible RTP pseudorotaxane in aqueous solutions.




Fig. 2 Schematic illustration of photo-controlled reversible RTP supramolecular assembly based on the azo- β -CD and α -BrNp in aqueous solutions.






Fig. 3 Schematic illustration of construction of the compound structure of photoresponsive supramolecular RTP gel based on β -CD and bromonaphthalene.




Fig. 4 Schematic illustration of construction of photocontrolled RTP supramolecular assembly in aqueous solution through poly-BrNpA/ γ -CD/poly-Azo system.

CBs are a class of water-soluble macrocycle compounds with different numbers of glycoluril units linked by methylene bridges that are able to form stable host-guest complexes with positively charged organic or inorganic guests in aqueous solution.^{11a, 11d, 12a, 23} The binding constant between CBs and guests is usually much larger than the corresponding value between CDs and guests due to the strong charge-dipole and hydrogen bonding interactions, as well as the hydrophobic/hydrophilic effect. This strong host-guest interaction can protect and immobilize the phosphor guest efficiently to improve their luminescent efficiency and lifetime in water. The RTP of quinoline and its derivatives could be induced through the formation of inclusion complexes with the different CBs in aqueous solution, and the effect of RTP was affected by the variation in heavy metal ions and pH.²⁴ In this system, the heavy atom effect of Ti^+ cation was stronger than that of the I^- anion, and the lower pH could enhance the RTP for the strong binding constant. The optical properties of inclusion complexes of different CBs with 2,4,6-triphenylpyrylium were studied in water, and the addition of CB[8] could produce strong RTP at 595 nm as a result of the CB carbonyl groups and the conformational immobilization inside the CB.²⁵ Tian and Ma also performed a series of studies on this feature. For example, a pH-controlled molecular shuttle in aqueous solution was prepared based on CB[7] and a 6-bromoisoquinoline derivative (Fig. 5).²⁶

In this shuttle, strong RTP was observed under the basic conditions, resulting from the strong electrostatic repulsion between the carboxyl anion of the deprotonated terminal carboxylic group and the high-density electron atmosphere of CB[7], leading to the encapsulation of the quinoline aromatic region in CB[7] for protecting the triplet state. Then, RTP was realized through the host-guest complex of CB[7] and 2-(4-aminobutyl)-6-bromo-1H-benzo[de]isoquinoline-1,3(2H)-dione (ANBrNpA) in water (Fig. 6).²⁷ The RTP was further enhanced by the association of 4-bromo-1,8-naphthalic anhydride polymer (poly-BrNpA) with CB[7]. Subsequently, a photoresponsive RTP system was constructed in aqueous solution by CB[7] and amphiphilic

6-bromoisoquinoline-functionalized alkoxyanthracene (AnBq).²⁸ In this system, the photoinduced electron transfer (PET) from anthracene to 6-bromoisoquinoline occurred for AnBq@CB[7], and therefore RTP was not observed. When AnBq@CB[7] underwent photolysis (Fig. 7), 6-bromoisoquinoline@CB[7], which lacked the anthracene group, had no intramolecular PET, resulting in phosphorescence. Recently, a visible light-excited water-soluble RTP was achieved via CB[8] and a double-sided modification of 4-(4-bromophenyl)-pyridine with the triazine core for the two-branched molecule TBP.²⁹ In this system, a 2:2 TBP-CB[8] quaternary complex was acquired as CB[8] was added into the TBP aqueous solution (Fig. 8). However, with the gradual increase in CB[8], the multicolor emission could be realized by controlling the different proportions of TBP quaternary complex, and the $(\text{TBP})_2@CB[8]_2$ assembly with yellow phosphorescence could also be regulated through the addition of a competitive guest (1-amino-3,5-dimethyladamantane hydrochloride). Meanwhile, a multicolor emission hydrogel was obtained via the mixture of $(\text{TBP})_2@CB[8]_2$ aqueous solution with different colors and an agarose gelator; finally, cell imaging was performed in the endosomes by taking the advantage of the visible light-excited RTP in water.



Fig. 5 Schematic illustration of the construction of a pH-controlled molecular shuttle in aqueous solution based on CB[7] and 6-bromoisoquinoline derivative.






Fig. 6 Schematic illustration of the construction of supramolecular aqueous RTP assembly based on CB[7] and ANBrNpA.




Fig. 7 Schematic illustration of construction of photoresponsive RTP supramolecular assembly by CB[7] and amphiphilic 6-bromoisoquinoline.




Fig. 8 Schematic illustration of construction of the TBP-CB[8] 2:2 quaternary RTP supramolecular assembly by CB[8] and TBP.

More recently, Liu and coworkers combined the 4-(4-bromophenyl)pyridinium bromide (BrBP) phosphor with a tumor-targeted hyaluronic acid (HA) polymer chain³⁰ subsequently assembled with cucurbituril (Fig. 9).³¹ BrBP, a type of organic phosphor, possessed a positive charge that could bind with CB.³² HA is a water-soluble, biocompatible, biodegradable polymer chain that can be used as a targeting reagent owing to specific binding with CD44 and RHAMM receptors that are overexpressed on the surface of cancer cells.³⁰ The encapsulation of CB[8] provided a rigid hydrophobic microenvironment to protect the phosphor guest from the collision of the triplet oxygen and reduce nonradiative decay as well as enhance ISC through the interaction of π - π stacking, halogen bonding, and multiple hydrogen bonding. The BrBP-modified HA formed small spherical aggregates. After

complexation with CB[8], larger aggregates were formed and resulted in purely organic RTP in water with a lifetime up to 4.73 ns (quantum yield 7.58%). In this system, the 1:2 strong inclusion complex of CB[8] with BrBP, as well as the interaction of CB[8]/BrBP complexes with HA, resulted in a strong charge-dipole, π - π stacking, hydrogen bonding interactions, and a hydrophobic effect, which prevented the influence of a quenching agent in aqueous solution, inhibited molecular rotation, and enhanced ISC. Furthermore, cell imaging was performed taking the advantages of cancer cell targeting property of HA via confocal laser scanning microscopy after incubation with the supramolecular polymer HA-BrBP/CB[8] and acquired optical signals of phosphorescence emission. The results in vitro revealed that the supramolecular polymer preferentially targeted the mitochondrion of tumor cells over normal cells and had negligible cytotoxic effects at the concentrations used for imaging. Meanwhile, the problem of UV excitation is expected to be solved by attempts to combine the supramolecular polymer with upconversion nanoparticles (UCNPs).

Meanwhile, a “two-end blocked” pseudorotaxane was constructed based on CB[8] and diethanolamine-modified bromopyridine salt (DA-PY), as shown in Fig. 10.³³ This special rotaxane structure could not only increase the phosphorescence emission in the water phase via reducing the nonradiative transition and enhancing the halogen bond interaction, but also realized the targeted imaging of cancer cell mitochondria. Additionally, supramolecular phosphorescent pins with afterglow luminescence and ultra-high quantum yield (99.38%) were designed through combining the assembly of alkyl-bridged phenylpyridinium salts and CB[8] into a rigid matrix (Fig. 11).³⁴ This high quantum yield arises from: 1) the suppressed nonradiative decay of the chromophore from the host-guest interaction of the chromophore with CB[8] and flexible chains; 2) the “one host and one guest” folding mode promoting the effective ICT for the increasing the rate of ISC; 3) the intramolecular halogen bond improving the rate of radiative decay. Furthermore, it was revealed that ICT and halogen bonding together promote RTP performance through the spectral comparison of reference compounds with only ICT structure or only halogen bond structure. However, there also exists a possibility other than the CT effect, where the main contribution comes from the heavy-atom induced RTP that is enhanced in a CB[8] cavity. Finally, this supramolecular pin was successfully used for phosphorescent imaging located in the mitochondria owing to the red-shift of the absorption.






Fig. 9 Schematic illustrating the self-assembly of ultralong water-soluble purely organic RTP supramolecular polymer by supramolecular and macromolecule effects.




Fig. 10 Schematic illustration of the construction of "two-end blocked" pseudorotaxane based on CB[8] and DA-PY.




Fig. 11 Schematic illustration of construction of supramolecular phosphorescent pins via the assembly of CB[8] and alkyl-bridged phenylpyridinium salts.

Subsequently, the emission of purely organic RTP could be controlled by introducing photoresponsive groups. Anthracene, a reversible response group that can perform photodimerization and photooxidation, is favored by scientists.³⁵ Moreover, the topological structure and fluorescence emission behavior could be controlled by modifying the anthracene groups on the π -aromatic cores, which prompted us to examine combining it with the phosphorescent groups to achieve the effective regulation of phosphorescent emission. Therefore, a bromophenylidinium salt was conjugated at the 2-position of anthracene (Fig. 12),³⁶ and the resultant compound displayed weak fluorescence emission at 502 nm. After adding CB[8], linear supramolecular aggregates were formed, and the fluorescence became stronger and red-shifted to 613 nm. In view of the photoresponse of anthracene, an illumination experiment showed that the morphology of the assembly was transformed from the linear head-to-tail inclusion complex into a head-to-head homoternary complex under 365 nm irradiation, accompanied by the conversion of fluorescence to phosphorescence (529 nm). Unexpectedly, this change in optical properties afforded two-organelle imaging from the nucleus to the lysosome in cell. In addition, diarylethylene derivatives are a class of molecules with photoisomerism and very good fatigue resistance.³⁷ Their configuration changes lead to different absorbances; this is conducive to the realization of optical property regulation. Therefore, the phosphorescent groups 4-(4-bromophenyl)-pyridinium and 4-(4-(5-methylthiophen-2-yl) phenyl) pyridinium were connected on both sides of diarylethylene through three carbon chains (Fig. 13).³⁸ In the presence of CB [7], due to the 1:1 bonding with the pyridine salt, the restriction of molecular motion is weak, and the emission of RTP cannot be achieved. In the presence of CB[8], owing to the strong 1:2 bonding with pyridine salt, π - π stacking could be enhanced to form a single-molecule dual-fold structure, and the RTP emission could be realized. Interestingly, the assembly emission overlapped well with the absorption of the near-infrared dye Cy5 (emission: 673 nm) to produce phosphorescent energy resonance transfer (PRET), and the process was regulated by the open-close ring of the diarylethylene. This light-controlled PRET assembly displayed good results in cell imaging.




Fig. 12 Schematic illustrating the construction of photoresponsive linear supramolecular aggregates based on the CB[8] and anthracene-modified bromophenylidinium salt.






Fig. 13 Schematic illustration of the construction of photocontrolled supramolecular phosphorescent energy transfer based on the diarylethene-modified bromophenylidinium salt.

The use of purely organic RTP and near-infrared light emitting dyes can solve the problem of shallow imaging depth caused by most RTP emission in the UV or visible range through energy transfer.^{74, 39} In a purely organic light-harvesting phosphorescence energy transfer system (Fig. 14) for a mitochondrially targeted imaging constructed from 4-(4-bromophenyl)-pyridine modified β -cyclodextrin (CD-PY), CB[8], rhodamine B (RhB), and adamantane-modified hyaluronic acid (HA-ADA), CD-PY@CB[8] acted as the phosphorescent donor, RhB acted as the phosphorescent acceptor, and HA-ADA acted as the cancer cell targeting agent.⁴⁰ Free CD-PY has no phosphorescence in water, and the manifested phosphorescence emission was observed at 510 nm as CB[8] was added to form host-guest inclusion complex. However, the PET process occurred when RhB was gradually added into the above system, and the phosphorescence of CD-PY@CB[8] decreased gradually at 510 nm with a new phosphorescence peak appearing at 590 nm. Interestingly, the addition of HA-ADA/CD into the above system not only further enhanced phosphorescence emission but also successfully presented mitochondrially targeted imaging property to cancer cells. For the supramolecular pin system, the emission of the folded assembly constructed by CB[8] and bromonaphthalene-modified methoxy pyridine changed from fluorescence at 425 nm to strong RTP at 530 nm (lifetime 130.2 μ s) as CB[8] enhanced the charge transfer interaction to promote ISC (Fig. 15).⁴¹ After the secondary assembly with amphiphilic calixarene (SC4AH), the G1/CB[8] assembly entered the hydrophobic region, and its phosphorescent property was further improved (luminescence intensity increased by 1.5 times, lifetime increased by 3.3 times). Moreover, a light-harvesting system from phosphorescence to near-infrared (NIR) fluorescence emission was realized using a ternary assembly G1@CB[8]@SC4AH as the donor and Nile red (emission: 635 nm) and Nile blue (emission: 675 nm) as the receptors (antenna effect: 352.9, 123.5; energy transfer efficiency 64.1%, 49.6%), respectively. This system was successfully applied to the NIR lysosome imaging of A549 cancer cells. Furthermore, by using this secondary assembly method, a two-staged phosphorescent energy transfer system was constructed (Fig. 16).⁴² First, di-

bromophthalimide derivative molecules were pre-assembled with CB[7], which promoted ISC to a certain extent and produced a weak phosphorescence at 505 nm when CB[7] was associated with the alkyl chain. After further assembly with amphiphilic SC4AH, the phosphorescence emission intensity of the assembly increased by 40 times, and the lifetime reached 1.13 ms. Taking the assembly as the donor and RhB or benzothiadiazole (DBT) as the receptor, efficient energy transfer from phosphorescence to fluorescence was achieved, with a transfer efficiency of 84.4% or 76.3%, and antenna effect of 289.4 or 119.5, respectively. Interestingly, after the addition of NIR dye Cy5 or Nile blue, which have better spectral overlap with RhB or DBT, the second-level phosphorescence to fluorescence energy transfer was realized, and its emission occurred over a wide range of emission from 425 nm to 800 nm. This secondary assembly was incubated with cells for 12 hours to explore the delay effect, and its emission at 505 nm, 585 nm, and 675 nm showed the corresponding time-resolved fluorescence signals after a delay of 50 μ s. Finally, the multicolor range imaging was realized in the lysosome of A549 cells.




Fig. 14 Schematic illustration of the construction of light-harvesting phosphorescence energy transfer supramolecular assembly via CD-PY, CB[8], RhB, and HA-ADA.




Fig. 15 Schematic illustration of the construction of light-harvesting phosphorescence energy transfer supramolecular assembly via SC4AH, CB[8], and bromonaphthalene-modified methoxy pyridine.






Fig. 16 Schematic illustration of construction of a two-staged light-harvesting phosphorescence energy transfer supramolecular assembly via SC4AH, CB[8], and di-bromophthalimide derivative.

Nanoassembly Confinement

The construction of nanoassemblies, including biocompatible and small size nanocrystals or aggregates, through hydrogen bonding, π - π stacking, and electrostatic interactions in aqueous media is an effective means to achieve the superior phosphorescence imaging of cells.^{43a, 44} The previously reported high-efficiency red phosphorescent compound C-Br was improved by the Liu group (Fig. 17),^{10a} and the carbazole and 4-bromobenzophenone groups were connected through an alkyl chain to give it a flexible conformation. The improved molecule can easily promote the formation of Br-H through a change in molecular structure stacking mode for increasing the intermolecular electronic coupling (IEC) and the heavy halogen atom effect, thus making the constructed nanocrystals display purely organic red phosphorescent emission in water, which is thus successfully applied in the afterglow imaging of cancer cells.




Fig. 17 Schematic illustration of the structure of phosphorescent compounds C-Br and C-C4-Br.

To avoid the problems of UV light-induced phototoxicity and lower light permeability to tissues and cells, three compounds with different substituents (H-NpCzBF₂, Br-NpCzBF₂, and I-NpCzBF₂) were prepared based on difluoroboron β -diketonate (BF₂bdk) and carbazole (Fig. 18),^{7j} in which BF₂bdk has two-photon character. In addition, the heavy atoms on the naphthalene molecule and the charge transfer and multiple bond interactions of the dimer could promote ISC and reduce nonradiative transitions for improving RTP

with NIR excitation. They then constructed H-NpCzBF₂, Br-NpCzBF₂, and I-NpCzBF₂ self-assembled dimeric nanoparticles in water by a nanoprecipitation method and achieved purely organic room temperature phosphorescence emission at approximately 630 nm by using two types of excitation (visible light at 470 nm or NIR light at 820 nm). Furthermore, the phosphorescence emission of three molecules was also observed in the cytoplasm of HeLa cells excited by visible or NIR light. However, the phosphorescence phenomenon based on carbazole derivatives was further confirmed by Liu and co-workers in 2021,⁸ⁱ in which the phosphorescence originates from the presence of a small number of carbazole isomers. Among them, two types of structures undergo charge transfer after photoexcitation to form a charge-separated state, and the charge-separated state is trapped and detrapped by defects, resulting in long-lived phosphorescence. However, to realize phosphorescence emission in aqueous solution, it is necessary to prepare nanoassemblies by π - π stacking, hydrogen bonding, and electrostatic interaction, which prevents nonradiative transition and the influence of free oxygen and other quenching groups on the high quantum yield. In addition, based on the sensitivity of phosphorescence to oxygen (O₂), the intracellular O₂ level could be read out accurately from the optical signals. The degree of hypoxia is an important parameter for the early diagnosis of cancer, cardiovascular disease, and stroke. For the quantification of molecular O₂, phosphorescence systems based on transition metal complexes are common; this is not only problematic because of the difficult synthesis, high cost, and cytotoxicity, but is also a challenge regarding stability. Subsequently, a 2-ureido-4-[1H]pyrimidinone (UPy)-functionalized bromine-substituted naphthalimide derivative was designed (BrNpA-UPy) (the UPy part is a quadruple hydrogen bond unit that can form a supramolecular polymer in chloroform), which could be further dimerized in chloroform with the bis-UPy modified benzyl (PH-bisUPy) through a microemulsion method, and then assembled in water to form supramolecular aggregate (Fig. 19).^{7b} Due to the interaction of quadruple hydrogen bonds and the heavy atom effect, the aggregate could emit a dual emission of 396 nm fluorescence and 570 nm phosphorescence at 348 nm under nitrogen conditions, revealing ratio change characteristics, notably that the phosphorescence disappears in air as it is sensitive to oxygen while the fluorescence emission part is unaffected. The aggregate could detect hypoxia in the cytoplasm of HeLa cancer cells by showing different emission colors in different oxygen concentration environments.




Fig. 18 Schematic illustration of the structure of the different substituents (H-NpCzBF₂, Br-NpCzBF₂, and I-NpCzBF₂) based on difluoroboron β -diketonate (BF₂bdk) and carbazole.






Fig. 19 Schematic illustration of the phosphorescence self-assembly of UPy-functionalized BrNpA-UPy.

Furthermore, water-soluble laponite (LP) clay nanoplates are an inorganic scaffold with a negatively charged surface and positively charged edge structure.³⁹ The positively charged edge can be pre-neutralized by the negatively charged polymer chain, which prevents the packing of LP. George et al. utilized the negatively charged surface of LP to assemble with positively charged bromine-substituted naphthalenediimine derivatives on both sides, thereby achieving deep red phosphorescence (emission = 615 nm) in water (Fig. 20).^{39a} Subsequently, they also reported a highly efficient light-harvesting system from phosphorescence to delayed fluorescence, which was assembled by heavy atom-substituted cationic phthalimide derivative/nanoclay hybrids (emission = 530 nm) with the negatively charged fluorescent dyes sulforhodamine G (emission = 560 nm) and sulforhodamine 101 (emission = 610 nm) with energy transfer efficiencies of 32% and 60% (D:A = 100:10), respectively (Fig. 21).^{39e}




Fig. 20 Schematic illustration of the proposed laponite ionic hybrid self-assembly.




Fig. 21 Schematic illustration of the LP-based phosphorescent harvesting assembly.

AIE is a phenomenon with high luminescence in an aggregated or confined state.⁴⁵⁻⁴⁶ Molecules with this property could overcome the quenching effect caused by aggregation. This property has a certain promoting effect on the realization of water phase phosphorescence. A hexathiobenzene persulfurated aromatic molecule was designed with six carboxylate groups on the periphery (Fig. 22),⁴⁷ which facilitated control of the ISC process through environmental changes to achieve RTP. The amphiphilic molecule 3 exhibits photoactivation and self-recovery of aggregation-induced phosphorescence (AIP) in water, and the reciprocation comes from the change in the degree of aggregation caused by the molecular conformational alteration and the dynamics of the dark lowest excited state. Similarly, the design of the two other reference molecules 1 and 2 confirmed this result. When an aqueous solution of molecule 3 was incubated with HeLa cancer cells, the photocontrollable phosphorescence effect was also achieved through rhythmic irradiation.




Fig. 22 Schematic illustration of the phosphorescent molecular structure.

Macromolecular Confinement



Influence factors of purely organic room temperature phosphorescence mainly have two aspects: one is the non-radiative transition, and the other is the quenching caused by the collision of oxygen.^{7f, 7h, 7i, 10b} In the solid state, the non-radiative transition can be minimized, and the collision with oxygen can also be reduced, so the design of solid-state purely organic room temperature phosphorescent molecules is easier to implement. In addition, amphiphiles are compounds or polymers with hydrophilic and hydrophobic fragments that can self-assemble to form well-defined structures in aqueous solutions, such as micelles and vesicles.⁴⁸ The hydrophobic drugs or molecules are encapsulated in hydrophobic fragments of amphiphiles through the non-covalent interactions (π - π stacking, hydrophobic forces and hydrogen-bonding), and the hydrophilic segment could not only increase the water-solubility and stability of the hydrophobic carrier but also improve the EPR effect in tumor part for enhancing the targeting.⁴⁹ Therefore, it is possible to combine the hydrophobic phosphorescent molecules with supramolecular amphiphilic system, so that their π - π stacking can be deposited in the hydrophobic region to prevent oxygen attacking, which may retain excellent phosphorescent property in solid state.

Frase et al.⁵⁰ reported the phosphorescence phenomena in water through nanoprecipitation method using iodide-substituted difluoroborondibenzoylmethane-poly(L-lactide) (BF2dbm(I)PLLA, 4) and methoxy-terminated poly(ethylene glycol)-b-poly(D-lactide) (mPEG-PDLA, 5) (Fig. 23). Compounds 1 and 2 could be assembled by stereo-complexation of hydrophobic PLA segments in DMF or mixed solvents (DMF/THF), with hydrophilic PEG chains outside. The constructed amphiphilic assembly dispersed in water, which could not only promote uptake of cancer cells by EPR effect but also realize phosphorescence/fluorescence colorimetric detection in tumor cells by using phosphorescence sensitivity to oxygen. The emission in the UV region is easily absorbed or scattered by absorbers in tissues, which limits biological imaging. Therefore, the conjugated structure of the dye was extended by substituting naphthalene to the benzene ring unit to red-shift the emission and reduce the scattering and absorption of the luminescence by the tissue structure; this consequently increased tissue penetration and enhanced the phosphorescence/fluorescence sensing and imaging efficiency of oxygen in living cells. (Fig. 24).⁵¹




Fig. 23 Schematic illustration of the structure of pegylated BNPs based on BF2dbm(I)PLLA (4) and mPEG-b-PDLA (5).




Fig. 24 Schematic illustration of the phosphorescent molecular structure and cell imaging.

Saponins, a class of amphiphilic sterols and triterpene glycosides with biological activity and biocompatibility that occurs naturally in plants, have good membrane permeabilization properties owing to the formation of transient pores by interacting with the cholesterol of the membrane above the critical micelle concentration (CMC).⁵² After combining RTP with aggregation-induced luminescence (AIE) molecules, the assembly had very good cell permeability and could enter cells very quickly (96 times faster than unencapsulated AIE molecules) (Fig. 25).⁴⁵ Furthermore, the long-lived purely organic RTP nanocrystals (NCs) 1-(dibenzo[b,d]furan-2-yl)phenylmethanone (BDBF) were encapsulated into nanorods to obtain an amphiphilic assembly that could quickly enter the cancer cells for long-lived luminescence imaging. The permeability of the unencapsulated monomer BDBF NCs in cells is very poor, but the amphiphilic assembly had better bleaching resistance compared with commercial dyes. Therefore, this amphiphilic assembly of purely organic room-temperature phosphorescence has very good prospects in biological applications.






Fig. 25 Schematic illustration of the cell permeabilization mechanism and AIE molecular structure.

PEG-*b*-PPG-*b*-PEG (F127), an amphiphilic block polymer composed of hydrophobic PPG and hydrophilic PEG, can form micelles through the aggregation of hydrophobic groups in aqueous solution.⁵³ Therefore, F127 can encapsulate the long-lived hydrophobic phosphorescent group in its hydrophobic portion in water. Six 10-phenyl-10H-phenothiazine-5,5-dioxide-based derivatives (i.e. CS-CH₃O, CS-CH₃, CSH, CS-Br, CS-Cl, and CS-F) were designed, in which different substituent groups were introduced to the 10-phenyl ring, including the electron-donating groups -CH₃O and -CH₃ and the electron-withdrawing groups -Br, -Cl, -F (Fig. 26).⁵⁴ The integration of theoretical calculations of electron cloud distribution and electric potential, etc., demonstrated the influence of substituent group changes on their phosphorescent properties. CS-F had the greater electron-withdrawing effect, which resulted in a shorter distance among the phenothiazine units and a stronger π - π interaction that could stabilize the excited triplet state efficiently to achieve ultralong RTP. Phosphorescence nanoparticles were then constructed with F127 and CS-F by the top-down approach in water to perform afterglow phosphorescent imaging of lymph nodes in living mice. Subsequently, six new compounds were constructed by replacing the benzene ring with a non-aromatic alkyl chain on the phenothiazine (Fig. 27).⁵⁵ Through experiments and theoretical calculations, the influence of alkyl chain length on phosphorescence was studied. It was found that the designed molecule had an odd/even effect; namely odd-numbered chains had stronger π - π interactions compared with even-numbered chains, leading to a longer phosphorescence effect. Among them, CS-C₃H₇, which has the longest lifetime RTP, was further combined with F127 and encapsulated in its hydrophobic region in water in a top-down manner. The phosphorescent imaging of human cancer cells in living mice has been achieved. In order to avoid problems with imaging

depth encountered by UV or visible light emission, the method of PRET was adopted to achieve afterglow purely organic RTP with NIR emission (Fig. 28).⁵⁶ *N,N*-bis(4-Methoxyphenyl)-3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) aniline (mTPA) emitted at 530 nm and silicon 2,3-naphthalocyanine bis(trihexylsilyloxy) (NCBS) emitted at 780 nm were also co-assembled with F127 in hydrophobic region in a top-down manner. Since the emission of mTPA and the absorption of NCBS partially overlapped, the NIR phosphorescent afterglow was realized in water by virtue of PRET. Furthermore, it was revealed that the effect of NIR afterglow imaging was shown by the aggregation in the lymph nodes of live mice.



Fig. 26 Schematic illustration of 10-phenyl-10H-phenothiazine-5,5-dioxide-based derivatives structures and the photograph of the corresponding molecular crystalline phosphorescence.




Fig. 27 Schematic illustration of molecular design strategy; and the preparation method of phosphorescent systems.






Fig. 28 Schematic illustration of PRET for in vivo imaging.

Outlook and perspectives

In conclusion, compared with fluorescence, aqueous RTP is becoming increasingly popular in the biochemical field owing to the advantages of larger Stokes shift, longer luminescence lifetime, and the involvement of triplet states, which can avoid the interference of background fluorescence or autofluorescence in complex biological systems. In particular, purely organic RTP, which has low cost, low toxicity, and easy modification, is favored by scientists. Additionally, a variety of weak interactions (π - π stacking, hydrogen bonding, and hydrophobicity) can provide a simple supramolecular strategy in water that can not only adjust the molecular configuration or electronic structure to change the properties of luminescent materials but also endow them dynamic intelligence. Supramolecular strategies may exhibit synergistic effects by combining two or more functional groups to avoid the disadvantages of a single component; this is widely used in biological and medical fields to meet current needs in the precision medicine field. Therefore, in this review, we systematically introduced the construction of purely organic RTP in the water phase by supramolecular means based on the application of cell imaging, including “macrocytic confinement,” “nanoassembly confinement,” and “macromolecule confinement.” It is worth noting that although purely organic RTP in water has been developed to a certain extent, the quantum yield and long lifetime in water still need to be improved, as the afterglow effect in water can better avoid the interference of background fluorescence in organisms. To achieve this, triplet excitons could be influenced by modulating the molecular structure, molecular aggregation morphology, or doping with other molecules, thereby improving the phosphorescence yield and lifetime of the molecule. However, UV light is highly phototoxic and has low penetration ability for cells and tissues, so the UV excitation as well as UV or visible light emission of phosphorescence in water is another shortcoming in the application of phosphorescence that is presently unresolved. However, the realization of purely organic RTP in water with visible light and even NIR light excitation as well as NIR light emission is the direction pursued

by researchers. In this regard, we believe that the excitation or emission redshift, or even a non-excitation light source, may be achieved by extending the degree of molecular conjugation, triplet-triplet energy transfer, or combining upconversion nanoparticles or chemiluminescence. Additionally, when combined with a variety of supramolecular noncovalent interactions through macrocyclic confinement, nanoconfinement and macromolecular confinement in water is expected to be an effective way to develop purely organic RTP with visible light to NIR excitation and NIR light emission, or even in the absence of an excitation light source, and thus to overcome this current barrier. Moreover, the stability of luminescent materials constructed by noncovalent interaction in living organisms still needs to be optimized and improved. Therefore, the development of purely organic RTP in water requires further effort. It is hoped that this review can provide some reference and inspiration for readers, as well as be a resource for the construction new aqueous purely organic RTP in biology.

Conflicts of interest

There are no conflicts to declare.

Author Contributions

Y.L. initiated the original inspiration for the manuscript. W.L.Z. wrote the manuscript. W.L. searched the literature and created the images. Y.L. and Y.C. edited and modified the manuscript.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (22101143, 22131008 and 21971127), the China Postdoctoral Science Foundation (2021M691661), the National Natural Science Foundation of Inner Mongolia (2021BS02014), and the Young Scientific and Technological Talents of Inner Mongolia (NJYT22050).

Notes and references

- (a) W. Li, S. Wu, X. Xu, J. Zhuang, H. Zhang, X. Zhang, C. Hu, B. Lei, C. F. Kaminski, Y. Liu, *Chem. Mater.* 2019, **31**, 9887-9894; (b) G. D. Luker, K. E. Luker, *J. Nucl. Med.* 2008, **49**, 1-4; (c) T. C. Voss, I. A. Demarco, R. N. Day, *BioTechniques* 2005, **38**, 413-424.
- (a) J. Chan, S. C. Dodani, C. J. Chang, *Nat. Chem.* 2012, **4**, 973-984; (b) D. Ding, K. Li, B. Liu, B. Z. Tang, *Acc. Chem. Res.* 2013, **46**, 2441-2453; (c) Y. Fan, P. Wang, Y. Lu, R. Wang, L. Zhou, X. Zheng, X. Li, J. A. Piper, F. Zhang, *Nat. Nanotechnol.* 2018, **13**, 941-946; (d) Kenry, Y. Duan, B. Liu, *Adv. Mater.* 2018, **30**, 1802394; (e) Z. Lei, F. Zhang, *Angew. Chem. Int. Ed.* 2021, **60**, 16294-16308; (f) J. Li, K. Pu, *Chem. Soc. Rev.* 2019, **48**, 38-71; (g) J. F. Lovell, T. W. B. Liu, J. Chen, G. Zheng, *Chem. Rev.* 2010, **110**, 2839-2857; (h) K. J. McHugh, L. Jing, A. M. Behrens, S. Jayawardena, W. Tang, M. Gao, R. Langer, A. Jaklenec, *Adv. Mater.* 2018, **30**, 1706356; (i) Q. Miao, C. Xie, X. Zhen, Y. Lyu, H. Duan, X. Liu, J. V. Jokerst, K. Pu, *Nat. Biotechnol.* 2017, **35**, 1102-1110; (j) G. M. van Dam, G. Themelis, L. M. A. Crane, N. J. Harlaar, R. G. Pleijhuis, W. Kelder, A. Sarantopoulos, J. S. de Jong, H. J. G. Arts, A. G. J. van der Zee, J. Bart, P. S. Low, V.



- Ntziachristos, *Nat. Med.* 2011, **17**, 1315-1319; (k) H. Wan, J. Yue, S. Zhu, T. Uno, X. Zhang, Q. Yang, K. Yu, G. Hong, J. Wang, L. Li, Z. Ma, H. Gao, Y. Zhong, J. Su, A. L. Antaris, Y. Xia, J. Luo, Y. Liang, H. Dai, *Nat. Commun.* 2018, **9**, 1171; (l) J. Zhang, X. Zhen, P. K. Upputuri, M. Pramanik, P. Chen, K. Pu, *Adv. Mater.* 2017, **29**, 1604764.
- 3 (a) J. Huang, J. Li, Y. Lyu, Q. Miao, K. Pu, *Nat. Mater.* 2019, **18**, 1133-1143; (b) B. Wang, Y. Wang, Y. Wang, Y. Zhao, C. Yang, Z. Zeng, S. Huan, G. Song, X. Zhang, *Anal. Chem.* 2020, **92**, 4154-4163; (c) K. Y. Zhang, Q. Yu, H. Wei, S. Liu, Q. Zhao, W. Huang, *Chem. Rev.* 2018, **118**, 1770-1839.
- 4 X. Zhen, Y. Tao, Z. An, P. Chen, C. Xu, R. Chen, W. Huang, K. Pu, *Adv. Mater.* 2017, **29**, 1606665.
- 5 (a) H. Liu, X. Zhang, B. Xing, P. Han, S. S. Gambhir, Z. Cheng, *Small* 2010, **6**, 1087-1091; (b) M.-K. So, C. Xu, A. M. Loening, S. S. Gambhir, J. Rao, *Nat. Biotechnol.* 2006, **24**, 339-343.
- 6 (a) Y. Chen, H. Zhang, Z. Lei, F. Zhang, *Small Struct.* 2020, **1**, 2000036; (b) O. Green, S. Gnaim, R. Blau, A. Eldar-Boock, R. Satchi-Fainaro, D. Shabat, *J. Am. Chem. Soc.* 2017, **139**, 13243-13248; (c) S. Gross, S. T. Gammon, B. L. Moss, D. Rauch, J. Harding, J. W. Heinecke, L. Ratner, D. Piwnica-Worms, *Nat. Med.* 2009, **15**, 455-461; (d) C. Zhang, K. Pu, *Small Struct.* 2020, **1**, 2000026.
- 7 (a) H. Gao, X. Ma, *Aggregate* 2021, **2**, e38; (b) F. Gu, X. Ma, *Chem. Eur. J.* 2022, e202104131; (c) L. Gu, H. Wu, H. Ma, W. Ye, W. Jia, H. Wang, H. Chen, N. Zhang, D. Wang, C. Qian, Z. An, W. Huang, Y. Zhao, *Nat. Commun.* 2020, **11**, 944; (d) L. Gu, W. Ye, X. Liang, A. Lv, H. Ma, M. Singh, W. Jia, Z. Shen, Y. Guo, Y. Gao, H. Chen, D. Wang, Y. Wu, J. Liu, H. Wang, Y.-X. Zheng, Z. An, W. Huang, Y. Zhao, *J. Am. Chem. Soc.* 2021, **143**, 18527-18535; (e) Z. He, H. Gao, S. Zhang, S. Zheng, Y. Wang, Z. Zhao, D. Ding, B. Yang, Y. Zhang, W. Z. Yuan, *Adv. Mater.* 2019, **31**, 1807222; (f) Z. Huang, X. Ma, *Cell Rep. Phys. Sci.* 2020, **1**, 100167; (g) X.-Q. Liu, K. Zhang, J.-F. Gao, Y.-Z. Chen, C.-H. Tung, L.-Z. Wu, *Angew. Chem. Int. Ed.* 2020, **59**, 23456-23460; (h) X. Ma, J. Wang, H. Tian, *Acc. Chem. Res.* 2019, **52**, 738-748; (i) X.-K. Ma, Y. Liu, *Acc. Chem. Res.* 2021, **54**, 3403-3414; (j) X.-F. Wang, H. Xiao, P.-Z. Chen, Q.-Z. Yang, B. Chen, C.-H. Tung, Y.-Z. Chen, L.-Z. Wu, *J. Am. Chem. Soc.* 2019, **141**, 5045-5050; (k) J. Yang, M. Fang, Z. Li, *Acc. Mater. Res.* 2021, **2**, 644-654; (l) Y. You, K. Huang, X. Liu, X. Pan, J. Zhi, Q. He, H. Shi, Z. An, X. Ma, W. Huang, *Small* 2020, **16**, 1906733; (m) X. Yu, W. Liang, Q. Huang, W. Wu, J. J. Chruma, C. Yang, *Chem. Commun.* 2019, **55**, 3156-3159; (n) T. Zhang, X. Ma, H. Wu, L. Zhu, Y. Zhao, H. Tian, *Angew. Chem. Int. Ed.* 2020, **59**, 11206-11216; (o) Y. Zhang, Y. Su, H. Wu, Z. Wang, C. Wang, Y. Zheng, X. Zheng, L. Gao, Q. Zhou, Y. Yang, X. Chen, C. Yang, Y. Zhao, *J. Am. Chem. Soc.* 2021, **143**, 13675-13685; (p) W. Zhao, Z. He, B. Z. Tang, *Nat. Rev. Mater.* 2020, **5**, 869-885; (q) W.-L. Zhou, W. Lin, Y. Chen, X.-Y. Dai, Z. Liu, Y. Liu, *Chem. Sci.* 2022, **13**, 573-579; (r) A. D. Nidhankar, Goudappagouda, V. C. Wakchaure, S. S. Babu, *Chem. Sci.* 2021, **12**, 4216-4236.
- 8 (a) A. Abdulkayum, J.-T. Chen, Q. Zhao, X.-P. Yan, *J. Am. Chem. Soc.* 2013, **135**, 14125-14133; (b) Z. An, C. Zheng, Y. Tao, R. Chen, H. Shi, T. Chen, Z. Wang, H. Li, R. Deng, X. Liu, W. Huang, *Nat. Mater.* 2015, **14**, 685-690; (c) R. E. Andrews, K. M. Shah, J. M. Wilkinson, A. Gartland, *Bone* 2011, **49**, 717-723; (d) O. Bolton, K. Lee, H.-J. Kim, K. Y. Lin, J. Kim, *Nat. Chem.* 2011, **3**, 205-210; (e) P. Collery, B. Keppler, C. Madoulet, B. Desoize, *Crit. Rev. Oncol. Hemat.* 2002, **42**, 283-296; (f) Q. le Masne de Chermont, C. Chanéac, J. Seguin, F. Pellé, S. Maîtrejean, J.-P. Jolivet, D. Gourier, M. Bessodes, D. Scherman, *Proc. Natl. Acad. Sci. U.S.A.* 2007, **104**, 9266; (g) Y. Li, M. Gecevicius, J. Qiu, *Chem. Soc. Rev.* 2016, **45**, 2090-2136; (h) T. Maldiney, A. Lecointre, B. Viana, A. Bessière, M. Bessodes, D. Gourier, C. Richard, D. Scherman, *J. Am. Chem. Soc.* 2011, **133**, 11810-11815; (i) C. Chen, Z. Chi, K. C. Chong, A. S. Batsanov, Z. Yang, Z. Mao, Z. Yang, B. Liu, *Nat. Mater.* 2021, **20**, 175-180.
- 9 X. Yang, D. Yan, *Chem. Sci.* 2016, **7**, 4519-4526.
- 10 (a) S. M. A. Fateminia, Z. Mao, S. Xu, Z. Yang, Z. Chi, B. Liu, *Angew. Chem. Int. Ed.* 2017, **56**, 12160-12164; (b) D. Li, J. Wang, X. Ma, *Adv. Opt. Mater.* 2018, **6**, 1800273.
- 11 (a) S. J. Barrow, S. Kaser, M. J. Rowland, J. del Barrio, O. A. Scherman, *Chem. Rev.* 2015, **115**, 12320-12406; (b) D. Dai, Z. Li, J. Yang, C. Wang, J.-R. Wu, Y. Wang, D. Zhang, Y.-W. Yang, *J. Am. Chem. Soc.* 2019, **141**, 4756-4763; (c) Z. Li, Y.-W. Yang, *Acc. Mater. Res.* 2021, **2**, 292-305; (d) G. Yu, K. Jie, F. Huang, *Chem. Rev.* 2015, **115**, 7240-7303; (e) W.-L. Zhou, Y. Chen, Y. Liu, *Acta Chim. Sinica* 2020, **78**, 1164-1176; (f) Y. Zhang, L. Wang, J. Wang, S. Xin, X. Sheng, *Chin. Chem. Lett.* 2021, **32**, 1902-1906.
- 12 (a) Y. Chen, F. Huang, Z.-T. Li, Y. Liu, *Sci. China Chem.* 2018, **61**, 979-992; (b) G. Crini, *Chem. Rev.* 2014, **114**, 10940-10975; (c) W.-F. Lai, A. L. Rogach, W.-T. Wong, *Chem. Soc. Rev.* 2017, **46**, 6379-6419; (d) J. Szejtli, *Chem. Rev.* 1998, **98**, 1743-1754; (e) W.-L. Zhou, Y. Chen, W. Lin, Y. Liu, *Chem. Commun.* 2021, **57**, 11443-11456.
- 13 N. J. Turro, J. D. Bolt, Y. Kuroda, I. Tabushi, *Photochem. Photobiol.* 1982, **35**, 69-72.
- 14 S. Scypinski, L. J. C. Love, *Anal. Chem.* 1984, **56**, 322-327.
- 15 A. Ponce, P. A. Wong, J. J. Way, D. G. Nocera, *J. Phys. Chem.* 1993, **97**, 11137-11142.
- 16 M. A. Mortellaro, W. K. Hartmann, D. G. Nocera, *Angew. Chem. Int. Ed.* 1996, **35**, 1945-1946.
- 17 Y.-S. Wei, W.-J. Jin, R.-H. Zhu, G.-W. Xing, C.-S. Liua, S.-S. Zhang, B.-L. Zhou, *Spectrochim. Acta A* 1996, **52**, 683-690.
- 18 Y.-X. Zhu, J.-H. Peng, Y. Zhang, *Anal. Chim. Acta* 2007, **583**, 364-369.
- 19 X. Ma, J. Cao, Q. Wang, H. Tian, *Chem. Commun.* 2011, **47**, 3559-3561.
- 20 J. Cao, X. Ma, M. Min, T. Cao, S. Wu, H. Tian, *Chem. Commun.* 2014, **50**, 3224-3226.
- 21 H. Chen, X. Ma, S. Wu, H. Tian, *Angew. Chem. Int. Ed.* 2014, **53**, 14149-14152.
- 22 H. Chen, L. Xu, X. Ma, H. Tian, *Polym. Chem.* 2016, **7**, 3989-3992.
- 23 (a) K. Kim, N. Selvapalam, Y. H. Ko, K. M. Park, D. Kim, J. Kim, *Chem. Soc. Rev.* 2007, **36**, 267-279; (b) X.-L. Ni, X. Xiao, H. Cong, Q.-J. Zhu, S.-F. Xue, Z. Tao, *Acc. Chem. Res.* 2014, **47**, 1386-1395; (c) J. Tian, L. Zhang, H. Wang, D.-W. Zhang, Z.-T. Li, *Supramol. Chem.* 2016, **28**, 769-783; (d) Z.-J. Yin, Z.-Q. Wu, F. Lin, Q.-Y. Qi, X.-N. Xu, X. Zhao, *Chin. Chem. Lett.* 2017, **28**, 1167-1171.
- 24 L. Mu, X.-B. Yang, S.-F. Xue, Q.-J. Zhu, Z. Tao, X. Zeng, *Anal. Chim. Acta* 2007, **597**, 90-96.
- 25 P. Montes-Navajas, H. Garcia, *J. Phys. Chem. C* 2010, **114**, 2034-2038.
- 26 Y. Gong, H. Chen, X. Ma, H. Tian, *ChemPhysChem* 2016, **17**, 1934-1938.
- 27 L. Xu, L. Zou, H. Chen, X. Ma, *Dyes Pigment.* 2017, **142**, 300-305.
- 28 T. Li, X. Ma, *Dyes Pigment.* 2018, **148**, 306-312.
- 29 J. Wang, Z. Huang, X. Ma, H. Tian, *Angew. Chem. Int. Ed.* 2020, **59**, 9928-9933.
- 30 (a) S. Jaracz, J. Chen, L. V. Kuznetsova, I. Ojima, *Bioorgan. Med. Chem.* 2005, **13**, 5043-5054; (b) Q. Yu, Y.-M. Zhang, Y.-H. Liu, X. Xu, Y. Liu, *Sci. Adv.* 2018, **4**, eaat2297.
- 31 W.-L. Zhou, Y. Chen, Q. Yu, H. Zhang, Z.-X. Liu, X.-Y. Dai, J.-J. Li, Y. Liu, *Nat. Commun.* 2020, **11**, 4655.
- 32 (a) Z.-Y. Zhang, Y. Chen, Y. Liu, *Angew. Chem. Int. Ed.* 2019, **58**, 6028-6032; (b) Z.-Y. Zhang, Y. Liu, *Chem. Sci.* 2019, **10**, 7773-7778.
- 33 X.-K. Ma, Y.-M. Zhang, Q. Yu, H. Zhang, Z. Zhang, Y. Liu, *Chem. Commun.* 2021, **57**, 1214-1217.
- 34 X.-K. Ma, W. Zhang, Z. Liu, H. Zhang, B. Zhang, Y. Liu, *Adv. Mater.* 2021, **33**, 2007476.



- 35 (a) G. Collet, T. Lathion, C. Besnard, C. Piguet, S. Petoud, *J. Am. Chem. Soc.* 2018, **140**, 10820-10828; (b) M. A. Filatov, S. Karuthedath, P. M. Polestshuk, H. Savoie, K. J. Flanagan, C. Sy, E. Sitte, M. Telitchko, F. Laquai, R. W. Boyle, M. O. Senge, *J. Am. Chem. Soc.* 2017, **139**, 6282-6285; (c) W. Zhou, Y. Chen, Q. Yu, P. Li, X. Chen, Y. Liu, *Chem. Sci.* 2019, **10**, 3346-3352; (d) Y. Zhou, H.-Y. Zhang, Z.-Y. Zhang, Y. Liu, *J. Am. Chem. Soc.* 2017, **139**, 7168-7171.
- 36 H.-J. Yu, Q. Zhou, X. Dai, F.-F. Shen, Y.-M. Zhang, X. Xu, Y. Liu, *J. Am. Chem. Soc.* 2021, **143**, 13887-13894.
- 37 (a) X. Dai, X. Dong, Z. Liu, G. Liu, Y. Liu, *Biomacromolecules* 2020, **21**, 5369-5379; (b) M. Irie, T. Fukaminato, K. Matsuda, S. Kobatake, *Chem. Rev.* 2014, **114**, 12174-12277; (c) Z. Li, G. Davidson-Rozenfeld, M. Vázquez-González, M. Fadeev, J. Zhang, H. Tian, I. Willner, *J. Am. Chem. Soc.* 2018, **140**, 17691-17701; (d) G. Liu, X. Xu, X. Dai, C. Jiang, Y. Zhou, L. Lu, Y. Liu, *Mater. Horiz.* 2021, **8**, 2494-2502; (e) D.-H. Qu, Q.-C. Wang, Q.-W. Zhang, X. Ma, H. Tian, *Chem. Rev.* 2015, **115**, 7543-7588.
- 38 C. Wang, X.-K. Ma, P. Guo, C. Jiang, Y.-H. Liu, G. Liu, X. Xu, Y. Liu, *Adv. Sci.* 2021, 2103041.
- 39 (a) S. Kuila, K. V. Rao, S. Garain, P. K. Samanta, S. Das, S. K. Pati, M. Eswaremoorthy, S. J. George, *Angew. Chem. Int. Ed.* 2018, **57**, 17115-17119; (b) S. Kuila, S. J. George, *Angew. Chem. Int. Ed.* 2020, **59**, 9393-9397; (c) L. Zhang, R. Zhang, P. Cui, W. Cao, F. Gao, *Chem. Commun.* 2013, **49**, 8102-8104; (d) B. Zhou, D. Yan, *Adv. Funct. Mater.* 2019, **29**, 1807599; (e) S. Garain, B. C. Garain, M. Eswaremoorthy, S. K. Pati, S. J. George, *Angew. Chem. Int. Ed.* 2021, **60**, 19720-19724; (f) X.-Y. Dai, Y.-Y. Hu, Y. Sun, M. Huo, X. Dong, Y. Liu, *Adv. Sci.* 2022, **9**, 2200524.
- 40 F.-F. Shen, Y. Chen, X. Dai, H.-Y. Zhang, B. Zhang, Y. Liu, Y. Liu, *Chem. Sci.* 2021, **12**, 1851-1857.
- 41 M. Huo, X.-Y. Dai, Y. Liu, *Small* 2022, 2104514.
- 42 M. Huo, X.-Y. Dai, Y. Liu, *Angew. Chem. Int. Ed.* 2021, **60**, 27171-27177.
- 43 (a) S. M. A. Fateminia, Z. Wang, C. C. Goh, P. N. Manghnani, W. Wu, D. Mao, L. G. Ng, Z. Zhao, B. Z. Tang, B. Liu, *Adv. Mater.* 2017, **29**, 1604100; (b) A. Pfister, G. Zhang, J. Zareno, A. F. Horwitz, C. L. Fraser, *ACS Nano* 2008, **2**, 1252-1258; (c) Y. S. Zhao, J. Xu, A. Peng, H. Fu, Y. Ma, L. Jiang, J. Yao, *Angew. Chem. Int. Ed.* 2008, **47**, 7301-7305.
- 44 X. Chen, C. Xu, T. Wang, C. Zhou, J. Du, Z. Wang, H. Xu, T. Xie, G. Bi, J. Jiang, X. Zhang, J. N. Demas, C. O. Trindle, Y. Luo, G. Zhang, *Angew. Chem. Int. Ed.* 2016, **55**, 9872-9876.
- 45 A. Nicol, R. T. K. Kwok, C. Chen, W. Zhao, M. Chen, J. Qu, B. Z. Tang, *J. Am. Chem. Soc.* 2017, **139**, 14792-14799.
- 46 (a) B. He, J. Huang, J. Zhang, H. H. Y. Sung, J. W. Y. Lam, Z. Zhang, S. Yan, D. Wang, J. Zhang, B. Z. Tang, *Angew. Chem. Int. Ed.* e202117709; (b) X. Liu, C. Zhu, B. Z. Tang, *Acc. Chem. Res.* 2022, **55**, 197-208; (c) J. Mei, N. L. C. Leung, R. T. K. Kwok, J. W. Y. Lam, B. Z. Tang, *Chem. Rev.* 2015, **115**, 11718-11940; (d) J. Wang, X. Gu, P. Zhang, X. Huang, X. Zheng, M. Chen, H. Feng, R. T. K. Kwok, J. W. Y. Lam, B. Z. Tang, *J. Am. Chem. Soc.* 2017, **139**, 16974-16979; (e) Z. Zhang, W. Yan, D. Dang, H. Zhang, J. Z. Sun, B. Z. Tang, *Cell Rep. Phys. Sci.* 2022, 100716.
- 47 X. Jia, C. Shao, X. Bai, Q. Zhou, B. Wu, L. Wang, B. Yue, H. Zhu, L. Zhu, *Proc. Natl. Acad. Sci. U.S.A.* 2019, **116**, 4816.
- 48 (a) H.-W. Tian, Y.-C. Liu, D.-S. Guo, *Mater. Chem. Front.* 2020, **4**, 46-98; (b) C. Wang, Z. Wang, X. Zhang, *Acc. Chem. Res.* 2012, **45**, 608-618; (c) X. Zhang, C. Wang, *Chem. Soc. Rev.* 2011, **40**, 94-101.
- 49 (a) L. Chen, Z. Xie, J. Hu, X. Chen, X. Jing, *J. Nanopart. Res.* 2007, **9**, 777-785; (b) M. R. Dreher, W. Liu, C. R. Michelich, M. W. Dewhirst, F. Yuan, A. Chilkoti, *J. Natl. Cancer Inst.* 2006, **98**, 335-344.
- 50 F. R. Kersey, G. Zhang, G. M. Palmer, M. W. Dewhirst, C. L. Fraser, *ACS Nano* 2010, **4**, 4989-4996.
- 51 C. A. DeRosa, J. Samonina-Kosicka, Z. Fan, H. C. Hendargo, D. H. Weitzel, G. M. Palmer, C. L. Fraser, *Macromolecules* 2015, **48**, 2967-2977.
- 52 (a) I. G. Barr, A. Sjölander, J. C. Cox, *Adv. Drug Deliv. Rev.* 1998, **32**, 247-271; (b) V. R. Dayeh, S. L. Chow, K. Schirmer, D. H. Lynn, N. C. Bols, *Ecotoxicol. Environ. Saf.* 2004, **57**, 375-382.
- 53 (a) P. Alexandridis, T. Alan Hatton, *Colloid. Surface. A* 1995, **96**, 1-46; (b) S. Kerkhofs, T. Willhammar, H. Van Den Noortgate, C. E. A. Kirschhock, E. Breynaert, G. Van Tendeloo, S. Bals, J. A. Martens, *Chem. Mater.* 2015, **27**, 5161-5169; (c) J. E. Matthew, Y. L. Nazario, S. C. Roberts, S. R. Bhatia, *Biomaterials* 2002, **23**, 4615-4619; (d) M. D. Shin, J. D. Hochberg, J. K. Pokorski, N. F. Steinmetz, *ACS Appl. Mater. Interfaces* 2021, **13**, 59618-59632.
- 54 J. Yang, X. Zhen, B. Wang, X. Gao, Z. Ren, J. Wang, Y. Xie, J. Li, Q. Peng, K. Pu, Z. Li, *Nat. Commun.* 2018, **9**, 840.
- 55 J. Yang, H. Gao, Y. Wang, Y. Yu, Y. Gong, M. Fang, D. Ding, W. Hu, B. Z. Tang, Z. Li, *Mater. Chem. Front.* 2019, **3**, 1391-1397.
- 56 Q. Dang, Y. Jiang, J. Wang, J. Wang, Q. Zhang, M. Zhang, S. Luo, Y. Xie, K. Pu, Q. Li, Z. Li, *Adv. Mater.* 2020, **32**, 2006752.

Supramolecular Assembly Confined Purely Organic Room-Temperature Phosphorescence and Its Biological Imaging

Wei-Lei Zhou^{a,b}, Wenjing Lin^a, Yong Chen^a and Yu Liu^{a*}

