


Cyclodextrin-Driven Movement of Cucurbit[7]uril

Yu Liu,* Xiao-Yun Li, Heng-Yi Zhang, Chun-Ju Li, and Fei Ding

Department of Chemistry, State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, People's Republic of China

yuliu@nankai.edu.cn

Received November 24, 2006



The movement of cucurbit[7]uril (CB[7]) driven by α -cyclodextrin (α -CD) is investigated by various experimental techniques including NMR, ESI-MS, UV-vis, and ITC. CB[7] can form stable pseudotaxanes with *N*-methyl-*N'*-octyl-4,4'-bipyridinium (MVO^{2+}) and *N,N'*-dioctyl-4,4'-bipyridinium (OV^{2+}) dication in aqueous solution. CB[7] shuttles between the octyl and bipyridinium moieties in MVO^{2+} , but docks at one of the octyl moieties in OV^{2+} . The addition of α -CD pushes CB[7] from the octyl moiety of MVO^{2+} or OV^{2+} to the bipyridinium moiety. Thermodynamically, the movement of CB[7] is mainly driven by exothermic enthalpy changes coming from the complexation of the octyl moiety of MVO^{2+} or OV^{2+} with α -CD.

Introduction

In the past few years interlocked supramolecules¹ have attracted a great deal of attention because of their interesting structures and potential application for supramolecular machines such as molecular switches,² molecular wires,^{3,4} and information

storage.⁵ Cucurbiturils (CBs) can form very stable host-guest complexes with positively charged molecules such as bipyridinium, ammonium, imidazolium, and coordinatively unsaturated metal ions through charge-dipole, hydrogen-bonding, and hydrophobic interactions.⁶ As another class of macrocycles contrasting with CBs, cyclodextrins (CDs) have been widely used as wheels for supramolecular assemblies for their remarkable properties of forming inclusion complexes with a variety

* To whom correspondence should be addressed. Fax: +86-22-23503625.

(1) (a) Lehn, J.-M. *Supramolecular Chemistry: Concepts and Perspectives*; Wiley-VCH: Weinheim, Germany, 1995. (b) Nepogodiev, S. A.; Stoddart, J. F. *Chem. Rev.* **1998**, *98*, 1959. (c) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2000**, *39*, 3348. (d) Collin, J.-P.; Dietrich-Buchecker, C.; Gavina, P.; Jimenez-Molero, M. C.; Sauvage, J.-P. *Acc. Chem. Res.* **2001**, *34*, 477. (e) Schalley, C. A.; Beizai, K.; Vögtle, F. *Acc. Chem. Res.* **2001**, *34*, 465. (f) Balzani, V.; Venturi, M.; Credi, A. *Molecular Devices and Machines*; Wiley-VCH: Weinheim, Germany, 2003.

(2) (a) Asakawa, M.; Higuchi, M.; Mattersteig, G.; Nakamura, T.; Pease, A. R.; Raymo, F. M.; Shimizu, T.; Stoddart, J. F. *Adv. Mater.* **2000**, *12*, 1099. (b) Bottari, G.; Leigh, D. A.; Pérez, E. *J. Am. Chem. Soc.* **2003**, *125*, 13360. (c) Feringa, B. L. *Acc. Chem. Res.* **2001**, *34*, 504. (d) Feringa, B. L. *Molecular Switches*; Wiley-VCH: Weinheim, Germany, 2001. (e) Leung, K. C.-F.; Mendes, P. M.; Magonov, S. N.; Northrop, B. H.; Kim, S.; Patel, K.; Flood, A. H.; Tseng, H.-R.; Stoddart, J. F. *J. Am. Chem. Soc.* **2006**, *128*, 10707. (f) Singhal, N. K.; Ramanujam, B.; Mariappanadar, V.; Rao, C. P. *Org. Lett.* **2006**, *8*, 3525.


(3) (a) Taylor, P. N.; O'Connell, M. J.; McNeill, L. A.; Hall, M. J.; Aplin, R. T.; Anderson, H. L. *Angew. Chem., Int. Ed.* **2000**, *39*, 3456. (b) Cacialli, F.; Wilson, J. S.; Michels, J. J.; Daniel, C.; Silva, C.; Friend, R. H.; Severin, N.; Samorì, P.; Rabe, J. P.; O'Connell, M. J.; Taylor, P. N.; Anderson, H. L. *Nat. Mater.* **2002**, *1*, 160. (c) Puigmarti-Luis, J.; Minoia, A.; Uji-i, H.; Rovira, C.; Cornil, J.; De Feyter, S.; Lazzaroni, R.; Amabilino, D. B. *J. Am. Chem. Soc.* **2006**, *128*, 12602. (d) Montes, V. A.; Perez-Bolivar, C.; Agarwal, N.; Shinar, J.; Anzenbacher, P., Jr. *J. Am. Chem. Soc.* **2006**, *128*, 12436.

(4) (a) Liu, Y.; You, C.-C.; Zhang, H.-Y.; Kang, S.-Z.; Zhu, C.-F.; Wang, C. *Nano Lett.* **2001**, *1*, 613. (b) Liu, Y.; Li, L.; Fan, Z.; Zhang, H.-Y.; Wu, X.; Liu, S.-X.; Guan, X.-D. *Nano Lett.* **2002**, *2*, 257-261. (c) Liu, Y.; Zhao, Y.-L.; Zhang, H.-Y.; Song, H.-B. *Angew. Chem., Int. Ed.* **2003**, *42*, 3260-3263. (d) Liu, Y.; Liang, P.; Chen, Y.; Zhang, Y.-M.; Zheng, J.-Y.; Yue, H. *Macromolecules* **2005**, *38*, 9095. (e) Zhao, Y.-L.; Zhang, H.-Y.; Guo, D.-S.; Liu, Y. *Chem. Mater.* **2006**, *18*, 4423.

(5) (a) Willner, I.; Pardo-Yissar, V.; Katz, E.; Ranjit, K. T. *J. Electroanal. Chem.* **2001**, *497*, 172. (b) Cavallini, M.; Biscarni, F.; LeWn, S.; Zerbetto, F.; Bottari, G.; Leigh, D. A. *Science* **2003**, *299*, 531.

(6) (a) Lee, J. W.; Samal, S.; Selvapalam, N.; Kim, H.-J.; Kim, K. *Acc. Chem. Res.* **2003**, *36*, 621. (b) Moon, K.; Kaifer, A. E. *Org. Lett.* **2004**, *6*, 185. (c) Sindelar, V.; Moon, K.; Kaifer, A. E. *Org. Lett.* **2004**, *6*, 2665. (d) Ong, W.; Kaifer, A. E. *J. Org. Chem.* **2004**, *69*, 1383. (e) Sindelar, V.; Cejas, M. A.; Raymo, F. M.; Kaifer, A. E. *New J. Chem.* **2005**, *29*, 289. (f) Ko, Y. H.; Kim, K.; Kang, J.-K.; Chun, H.; Lee, J. W.; Sakamoto, S.; Yamaguchi, K.; Fettinger, J. C.; Kim, K. *J. Am. Chem. Soc.* **2004**, *126*, 1932. (g) Kim, J.; Jung, I.-S.; Kim, S.-Y.; Lee, E.; Kang, J.-K.; Sakamoto, S.; Yamaguchi, K.; Kim, K. *J. Am. Chem. Soc.* **2000**, *122*, 540. (h) Mohanty, J.; Bhasikuttan, A. C.; Nau, W. M.; Pal, H. *J. Phys. Chem. B* **2006**, *110*, 5132. (i) Ong, W.; Kaifer, A. E. *Organometallics* **2003**, *22*, 4181. (j) Kim, K. *Chem. Soc. Rev.* **2002**, *31*, 96. (k) Rekharsky, M. V.; Yamamura, H.; Inoue, C.; Kawai, M.; Osaka, I.; Arakawa, R.; Shiba, K.; Sato, A.; Ko, Y. H.; Selvapalam, N.; Kim, K.; Inoue, Y. *J. Am. Chem. Soc.* **2006**, *128*, 14871.

CHART 1. Structures of CB[7] and the Viologen Guests



of small molecules or ions of appropriate size.⁷ 4,4'-Bipyridinium dication, commonly known as viologen, has been extensively utilized as a component of macrorings or axles due to its unique redox character and ability as an electron acceptor.^{7b,8} Recently, Inoue and co-workers studied the 1:1:1 CB[6]–CD–dihexylammonium complex that mainly depended on the supramolecular positive cooperativity.⁹ The pH-controlled and electrochemically switchable pseudorotaxanes based on CB[7] and viologens (Chart 1) have also been reported by Kaifer's group.¹⁰ Kim and Yui et al. studied pH-responsive movement of CB[7] in polypseudorotaxane containing dimethyl- β -CD and CB[7].¹¹ However, the movement of one macrocyclic molecule controlled by another macrocyclic molecule is not involved. Therefore, we are motivated to investigate in detail the movement of CB[7] on the axles of the linear viologens driven by α -CD.

It is documented that the binding interactions between symmetrical viologens and CB[7] may lead to two different types of complexes:^{6b} (a) *N,N'*-dimethyl-4,4'-bipyridinium (MV^{2+}) (Chart 1) or *N,N'*-diethyl-4,4'-bipyridinium (EV^{2+}) forms stable pseudorotaxane complexes with CB[7], in which CB[7] binds at the viologen nucleus; (b) *N,N'*-dibutyl-4,4'-bipyridinium (BV^{2+}) (Chart 1) or other viologens with longer aliphatic *N*-substituents favor the formation of external complexes with CB[7], in which CB[7] binds at the external docking points of viologens. In addition, α -CD hardly forms any complexes with dicationic guests such as viologens.¹² On the contrary, its cavity can include small guest molecules possessing straight aliphatic chains.¹³ In the text, we respectively chose asymmetric *N*-methyl-*N'*-octyl-4,4'-bipyridinium (MVO^{2+}) and symmetrical MV^{2+} , BV^{2+} , *N,N'*-dioctyl-4,4'-bipyridinium (OV^{2+}), and *N,N'*-dibenzyl-4,4'-bipyridinium (BnV^{2+}) as axles (Chart 1) and CB[7] and α -CD as wheels to build pseudorotaxanes. Our particular interest is to investigate how CB[7] moves on these viologen axles upon addition of α -CD.




FIGURE 1. ^1H NMR spectra (300 MHz, 0.2 M NaCl– D_2O , 298 K) of (A) MVO^{2+} , (B) MVO^{2+} + 0.3 equiv of CB[7], (C) MVO^{2+} + 1.1 equiv of CB[7], (D) MVO^{2+} + 1.1 equiv of CB[7] + 1.2 equiv of α -CD, (E) MVO^{2+} + 1.1 equiv of CB[7] + 3.0 equiv of α -CD, (F) MVO^{2+} + 1.1 equiv of CB[7] + 10.0 equiv of α -CD, and (G) MVO^{2+} + 10.0 equiv of α -CD.

Results and Discussion

^1H NMR Spectra. The formation of the 1:1 or 1:2 complexes for viologen guests with CB[7] and the 1:1:1 or 1:1:2 complexes for MVO^{2+} or OV^{2+} with CB[7] and α -CD can be conveniently monitored by ^1H NMR spectroscopy. We started our investigation with MVO^{2+} , CB[7], and α -CD. Figure 1A shows the ^1H NMR spectra of MVO^{2+} in 0.2 M NaCl– D_2O . Upon addition of 0.3 equiv of CB[7], the signals of aromatic protons are almost unchanged and those of the alkyl protons are broadened (Figure 1B). In the presence of 1.1 equiv of CB[7], the resonances of both the aromatic protons (β , β' , and α') and the alkyl protons shift to higher field (Figure 1C). This reveals that CB[7] is equilibrating between the bipyridinium and the octyl moieties in $[\text{MVO}\cdot\text{CB}[7]]^{2+}$, but mainly locates at the octyl moiety. Upon addition of 1.2 equiv of α -CD to the above mixed solution, the signals of β, β' aromatic protons exhibit a major upfield shift, and simultaneously those of octyl protons exhibit a downfield shift (Figure 1D). These observations indicate that CB[7] should bind at the bipyridinium moiety in the ternary complex $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$. The almost similar information of the chemical shifts of the aromatic and octyl protons in Figure 1E,F means that 3.0 equiv of α -CD can make all CB[7] locate at the bipyridinium moiety of MVO^{2+} . In the control experiment (Figure 1G), the addition of α -CD gives the signals of the octyl protons a minor downfield shift, but does not affect the chemical shift of the aromatic protons of MVO^{2+} , suggesting that α -CD just binds at the octyl moiety of MVO^{2+} . These combined observations confirm that CB[7] locates at the bipyridinium moiety of MVO^{2+} along with the addition of α -CD instead of shuttling between the octyl and bipyridinium moieties in $[\text{MVO}\cdot\text{CB}[7]]^{2+}$. The formation of the binary $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ and the ternary $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$ host–guest complexes is also evidenced by ESI-MS. The peaks at 723.8 and 1209.5 are

(7) (a) Harada, A. *Acc. Chem. Res.* **2001**, *34*, 456. (b) Park, J. W.; Song, H. J. *Org. Lett.* **2004**, *6*, 4869. (c) Anderson, S.; Claridge, T. D. M.; Anderson, H. L. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1310. (d) Lim, C. W.; Sakamoto, S.; Yamaguchi, K.; Hong, J.-I. *Org. Lett.* **2004**, *6*, 1079. (e) Miyachi, M.; Hoshino, T.; Yamaguchi, H.; Kamitori, S.; Harada, A. *J. Am. Chem. Soc.* **2005**, *127*, 2034. (f) Mezzina, E.; Fani, M.; Ferroni, F.; Franchi, P.; Menna, M.; Lucarini, M. *J. Org. Chem.* **2006**, *71*, 3773.

(8) Alvaro, M.; Ferrer, B.; Garcia, H.; Palomares, E. J.; Balzani, V.; Credi, A.; Venturi, M.; Stoddart, J. F.; Wenger, S. *J. Phys. Chem. B* **2003**, *107*, 14319.

(9) Rekharsky, M. V.; Yamamura, H.; Kawai, M.; Osaka, I.; Arakawa, R.; Sato, A.; Ko, Y. H.; Selvapalam, N.; Kim, K.; Inoue, Y. *Org. Lett.* **2006**, *8*, 815.

(10) (a) Sindelar, V.; Silvi, S.; Kaifer, A. E. *Chem. Commun.* **2006**, 2185. (b) Sobransingh, D.; Kaifer, A. E. *Org. Lett.* **2006**, *8*, 3247.

(11) Ooya, T.; Inoue, D.; Choi, S. H.; Kobayashi, Y.; Loethen, S.; Thompson, D. H.; Ko, Y. H.; Kim, K.; Yui, N. *Org. Lett.* **2006**, *8*, 3159.

(12) (a) Matsue, T.; Kato, T.; Akiba, U.; Osa, T. *Chem. Lett.* **1985**, *14*, 1825. (b) Mirzozian, A.; Kaifer, A. E. *Chem.–Eur. J.* **1997**, *3*, 1052.

(13) Rekharsky, M. V.; Inoue, Y. *Chem. Rev.* **1998**, *98*, 1875.




FIGURE 2. ESI-MS spectrum of a solution containing MVO^{2+} (1.0 equiv), $\text{CB}[7]$ (1.2 equiv), and $\alpha\text{-CD}$ (2.0 equiv) in water.




FIGURE 3. ^1H NMR spectra (300 MHz, 0.2 M $\text{NaCl-D}_2\text{O}$, 298 K) of (A) OV^{2+} , (B) OV^{2+} + 0.5 equiv of $\text{CB}[7]$, (C) OV^{2+} + 1.2 equiv of $\text{CB}[7]$, and (D) OV^{2+} + 2.4 equiv of $\text{CB}[7]$.

assigned to $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ and $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$, respectively, as shown in Figure 2.

We also chose the symmetrical OV^{2+} guest to investigate the movement of $\text{CB}[7]$ on viologen driven by $\alpha\text{-CD}$. As shown in Figure 3B, the presence of 0.5 equiv of $\text{CB}[7]$ splits every aliphatic proton signal of OV^{2+} into two signals. In the presence of 2.4 equiv of $\text{CB}[7]$, the aromatic proton signals are hardly changed, and the octyl ones obviously shift upfield (Figure 3D). These observations suggest that a 1:2 complex of $[\text{OV}\cdot 2\text{CB}[7]]^{2+}$ is formed, and the two $\text{CB}[7]$'s are respectively docked at the two octyl moieties in $[\text{OV}\cdot 2\text{CB}[7]]^{2+}$. The further NMR experiments of the addition of $\alpha\text{-CD}$ to $[\text{OV}\cdot 2\text{CB}[7]]^{2+}$ reveal that $\alpha\text{-CD}$ cannot change the binding sites of $\text{CB}[7]$ at OV^{2+} .¹⁴

The presence of 1.2 equiv of $\text{CB}[7]$ makes the octyl signals split into three portions (Figure 4A). They should correspond with the 1:1 complex of $[\text{OV}\cdot\text{CB}[7]]^{2+}$ and 1:2 complex of $[\text{OV}\cdot$




FIGURE 4. ^1H NMR spectra (300 MHz, 0.2 M $\text{NaCl-D}_2\text{O}$, 298 K) of (A) OV^{2+} + 1.2 equiv of $\text{CB}[7]$, (B) OV^{2+} + 1.2 equiv of $\text{CB}[7]$ + 1.2 equiv of $\alpha\text{-CD}$, (C) OV^{2+} + 1.2 equiv of $\text{CB}[7]$ + 4.0 equiv of $\alpha\text{-CD}$, (D) OV^{2+} + 1.2 equiv of $\text{CB}[7]$ + 8.0 equiv of $\alpha\text{-CD}$, (E) OV^{2+} + 1.2 equiv of $\text{CB}[7]$ + 12.0 equiv of $\alpha\text{-CD}$, and (F) OV^{2+} + 10.0 equiv of $\alpha\text{-CD}$.

$2\text{CB}[7]]^{2+}$. From their integral areas, the dominating component between them is $[\text{OV}\cdot\text{CB}[7]]^{2+}$. Furthermore, upon stepwise addition of $\alpha\text{-CD}$ to the mixed solution of OV^{2+} and 1.2 equiv of $\text{CB}[7]$, upfield displacement of the β aromatic protons is observed, and the signals of the alkyl protons are displaced downfield and broadened (from part B to part E of Figure 4). Upon addition of 12.0 equiv of $\alpha\text{-CD}$, the resonance signals of the β aromatic protons shift upfield from 8.42 to 7.45 ppm, and those of the alkyl protons are similar to the signals of OV^{2+} in the presence of 10.0 equiv of $\alpha\text{-CD}$, as shown in Figure 4F. This pattern of complexation-induced shifts could be ascribed to $\text{CB}[7]$ binding at the viologen nucleus, while $\alpha\text{-CD}$ includes the octyl chains in $[\text{OV}\cdot\text{CB}[7]\cdot 2\alpha\text{-CD}]^{2+}$. The $[\text{OV}\cdot\text{CB}[7]]^{2+}$, $[\text{OV}\cdot 2\text{CB}[7]]^{2+}$, $[\text{OV}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$, and $[\text{OV}\cdot\text{CB}[7]\cdot 2\alpha\text{-CD}]^{2+}$ complexes are unambiguously detected by ESI-MS measurements.¹⁴ It should be noted that OV^{2+} possesses two long alkyl chains, so a small quantity of $\alpha\text{-CD}$ cannot move $\text{CB}[7]$ to the viologen nucleus. In light of the above NMR and MS experiments, we deduced that the movement of $\text{CB}[7]$ is as follows: $\alpha\text{-CD}$ first binds at the free octyl moiety of $[\text{OV}\cdot\text{CB}[7]]^{2+}$, and then the binding of the second $\alpha\text{-CD}$ with another octyl chain of OV^{2+} forces $\text{CB}[7]$ to move from the octyl moiety to the bipyridinium moiety. In the process, both $\alpha\text{-CD}$ s play the roles of stoppers.

For comparison purposes, we investigated the binding interactions of other viologens (BV^{2+} , BnV^{2+} , and MV^{2+}) with $\text{CB}[7]$ and $\alpha\text{-CD}$ by NMR spectroscopy.¹⁴ BV^{2+} and BnV^{2+} can form external inclusion complexes with $\text{CB}[7]$ as OV^{2+} , in which the viologen nucleus is not engulfed by the $\text{CB}[7]$.^{6b,c} In sharp contrast, the addition of abundant $\alpha\text{-CD}$ cannot push $\text{CB}[7]$ to the viologen nucleus. One possible reason is that both external butyl and benzyl groups are too short to be included by $\alpha\text{-CD}$. For the case of MV^{2+} , $\text{CB}[7]$ binds at the viologen nucleus directly,^{6b,15} and the addition of $\alpha\text{-CD}$ cannot change the binding mode of $[\text{MV}\cdot\text{CB}[7]]^{2+}$.

(14) See the the Supporting Information.

(15) Kim, H.-J.; Jeon, W. S.; Ko, Y. H.; Kim, K. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 5007.




FIGURE 5. UV-vis spectra of 50.0 μM $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ in the presence of increasing concentrations (0–8.0 mM, in the direction of the arrow) of $\alpha\text{-CD}$ in Tris buffer (pH 7.4) at 298 K.

UV-Vis Spectra. UV-vis spectroscopic measurements were widely utilized to measure the binding constants between viologens and CB[7] because the molar absorptivity coefficients of their electronic absorption bands at ca. 260 nm were depressed when the viologen nucleus was included inside the cavity of CB[7]. In our experiments, the main parameters (λ_{max} and ϵ) of the UV absorption band of MVO^{2+} and OV^{2+} were not changed significantly by the increase of CB[7] or $\alpha\text{-CD}$. Interestingly, the addition of $\alpha\text{-CD}$ showed a small hypochromic shift in the absorption band of MVO^{2+} in the presence of 1.0 equiv of CB[7] as shown in Figure 5, indicating the anticipated locomotion of CB[7] from the octyl moiety to the viologen nucleus. Analogous results were also obtained from the spectra of the addition of $\alpha\text{-CD}$ to the $[\text{OV}\cdot\text{CB}[7]]^{2+}$ aqueous Tris buffer solution.¹⁴ However, the main parameters (λ_{max} and ϵ) of $[\text{BnV}\cdot\text{CB}[7]]^{2+}$, $[\text{BV}\cdot\text{CB}[7]]^{2+}$, or $[\text{MV}\cdot\text{CB}[7]]^{2+}$ were unchanged upon addition of $\alpha\text{-CD}$. These results from the UV-vis spectra were in excellent agreement with the NMR experiments.

Microcalorimetric Titration. To investigate quantitatively the molecular recognition behavior of CB[7], $\alpha\text{-CD}$ with viologens, and their thermodynamic origins for the movement of CB[7] on viologens driven by $\alpha\text{-CD}$, the isothermal titration calorimetry (ITC) experiments were performed in aqueous Tris buffers (pH 7.4). It should be mentioned that the protonated form of Tris can compete to some extent for CB[7] with the viologen dications.^{6d} As can be seen from Table 1, the K_S value for the complexation of MVO^{2+} with CB[7] is 2 orders of magnitude larger than that with $\alpha\text{-CD}$, leading to a large free energy change ($\Delta\Delta G^\circ = -11.8 \text{ kJ}\cdot\text{mol}^{-1}$). The favorable free




FIGURE 6. Partial ^1H ROESY spectrum of a mixture of MVO^{2+} (5 mmol/L), CB[7] (5 mmol/L), and $\alpha\text{-CD}$ (6 mmol/L) in D_2O at 298 K with a mixing time of 250 ms.

energy change results from the contributions of both the enthalpy term originating from ion-dipole interactions between the N^+ groups of MVO^{2+} and the carbonyl groups of CB[7] ($\Delta\Delta H^\circ = -4.7 \text{ kJ}\cdot\text{mol}^{-1}$) and the entropy term from the desolvation effect of the N^+ groups ($\Delta T\Delta S^\circ = 7.1 \text{ kJ}\cdot\text{mol}^{-1}$). It is noted that the complex $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ can interact further with $\alpha\text{-CD}$ to give the 1:1:1 $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$ complex. If CB[7] exclusively locates at the bipyridinium moiety of MVO^{2+} originally, the thermodynamic parameters of $\alpha\text{-CD}$ with $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ should be similar to those of the complexation of $\alpha\text{-CD}$ with MVO^{2+} . However, as shown in Table 1, the ΔG° value of the reaction is almost the same as that for complexation of MVO^{2+} with $\alpha\text{-CD}$, but their enthalpy changes and entropy changes are distinctly different: the complexation of $\alpha\text{-CD}$ with MVO^{2+} mostly comes from the contribution of the enthalpy, while that with $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ is attributed to both favored enthalpy change and entropy change. It should be mentioned that the favored enthalpy change for the formation of $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$ comes from not only the complexation of the octyl moiety with $\alpha\text{-CD}$ but also the hydrogen-bonding interactions

TABLE 1. Stability Constant (K_S/M^{-1}) and Standard Free Energy ($\Delta G^\circ/\text{kJ}\cdot\text{mol}^{-1}$), Enthalpy ($\Delta H^\circ/\text{kJ}\cdot\text{mol}^{-1}$), and Entropy ($T\Delta S^\circ/\text{kJ}\cdot\text{mol}^{-1}$) Changes for 1:1 Inclusion Complexation of MVO^{2+} with CB[7] or $\alpha\text{-CD}$ and $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ with $\alpha\text{-CD}$ and 1:2 Inclusion Complexation of OV^{2+} with CB[7] or $\alpha\text{-CD}$ and $[\text{OV}\cdot\text{CB}[7]]^{2+}$ with $\alpha\text{-CD}$ in Tris Buffer (pH 7.4) at 298 K

complexation reaction	K_S	$-\Delta G^\circ$	$-\Delta H^\circ$	$T\Delta S^\circ$
$\text{MVO}^{2+} + \text{CB}[7] \rightleftharpoons [\text{MVO}\cdot\text{CB}[7]]^{2+a}$	338000 ± 8000	31.56 ± 0.06	24.1 ± 0.2	7.5 ± 0.2
$\text{MVO}^{2+} + \alpha\text{-CD} \rightleftharpoons [\text{MVO}\cdot\alpha\text{-CD}]^{2+a}$	2940 ± 40	19.80 ± 0.04	19.4 ± 0.2	0.4 ± 0.2
$[\text{MVO}\cdot\text{CB}[7]]^{2+} + \alpha\text{-CD} \rightleftharpoons [\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+a}$	2180 ± 20	19.06 ± 0.03	8.0 ± 0.2	11.1 ± 0.2
$\text{OV}^{2+} + 2\text{CB}[7] \rightleftharpoons [\text{OV}\cdot 2\text{CB}[7]]^{2+a}$	516000 ± 20000	32.60 ± 0.09	21.7 ± 0.2	10.9 ± 0.3
$\text{OV}^{2+} + 2\alpha\text{-CD} \rightleftharpoons [\text{OV}\cdot 2\alpha\text{-CD}]^{2+a}$	2360 ± 24	19.24 ± 0.03	22.25 ± 0.06	-3.08 ± 0.09
$[\text{OV}\cdot\text{CB}[7]]^{2+} + \alpha\text{-CD} \rightleftharpoons [\text{OV}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+b}$	5640 ± 250	21.4 ± 0.1	14.8 ± 0.2	6.6 ± 0.1
$[\text{OV}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+} + \alpha\text{-CD} \rightleftharpoons [\text{OV}\cdot\text{CB}[7]\cdot 2\alpha\text{-CD}]^{2+b}$	626 ± 16	16.0 ± 0.1	19.2 ± 0.4	-3.3 ± 0.5

^a The net reaction heat in this run was calculated by the “one set of binding sites” model in ORIGIN software. ^b The net reaction heat in this run was calculated by the “1:2 sequential binding sites” model in ORIGIN software.




FIGURE 7. Possible structures of (a) $[\text{MVO}\cdot\text{CB}[7]]^{2+}$, (b) $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$, (c) $[\text{OV}\cdot\text{CB}[7]]^{2+}$, and (d) $[\text{OV}\cdot\text{CB}[7]\cdot 2\alpha\text{-CD}]^{2+}$ constructed according to MD simulation.

between CB[7] and α -CD. These observations suggest that CB[7] in the complex $[\text{MVO}\cdot\text{CB}[7]]^{2+}$ should mainly locate at the octyl moiety of MVO^{2+} , and the addition of α -CD pushes CB[7] from the octyl moiety to the bipyridinium moiety.

For OV^{2+} possessing two octyl chains, the 2:1 and 2:1:1 inclusion complexes are respectively formed between CB[7] (α -CD) and OV^{2+} and between α -CD and $[\text{OV}\cdot\text{CB}[7]]^{2+}$. The binding constants (K_S) and the relevant thermodynamic parameters obtained are shown in Table 1. The K_S values for the complexation of CB[7] or α -CD with OV^{2+} approximate those with MVO^{2+} , but their enthalpy changes and entropy changes are different. In comparison to the latter, the complexation of CB[7] with OV^{2+} gives a favorable entropy change ($\Delta T\Delta S^\circ = 3.4 \text{ kJ}\cdot\text{mol}^{-1}$) and a less unfavorable enthalpy change ($\Delta\Delta H^\circ = 2.4 \text{ kJ}\cdot\text{mol}^{-1}$). Contrarily, the complexation of α -CD and OV^{2+} is enthalpically favorable ($\Delta\Delta H^\circ = -2.8 \text{ kJ}\cdot\text{mol}^{-1}$) and entropically unfavorable ($\Delta T\Delta S^\circ = -3.5 \text{ kJ}\cdot\text{mol}^{-1}$). Further, the binding constant for the formation of the 1:1 complex of α -CD and $[\text{OV}\cdot\text{CB}[7]]^{2+}$ ($5640 \pm 250 \text{ M}^{-1}$) is larger than that of α -CD and OV^{2+} ($2360 \pm 24 \text{ M}^{-1}$). This should be ascribed to the OV^{2+} charge density decreasing with the binding of CB[7] and resulting in α -CD easily binding at the octyl moiety.

The above NMR experiments have clarified that CB[7] just binds at the octyl moiety of $[\text{OV}\cdot\text{CB}[7]]^{2+}$, while CB[7] does at the viologen nucleus of MV^{2+} . The binding constant of CB[7] with OV^{2+} is slightly larger than that with MV^{2+} ($K_S = 2.24 \times 10^5 \text{ M}^{-1}$,^{6d} $2.0 \times 10^5 \text{ M}^{-1}$,¹⁵ and $1.03 \times 10^5 \text{ M}^{-1}$,¹⁶). That is to say, the affinity of CB[7] toward the octyl moiety of OV^{2+} should be slightly stronger than that toward its bipyridinium moiety. Therefore, it is a natural process that the weaker binding of α -CD with OV^{2+} can compete with very strong binding of CB[7] to force it to move from the octyl moiety to the bipyridinium moiety.

Two-Dimensional NMR Experiments. To investigate the interactions among MVO^{2+} , CB[7], and α -CD, 2D NMR experiments were employed upon addition of 1.2 equiv of α -CD to a solution of $[\text{MVO}\cdot\text{CB}[7]]^{2+}$. As shown in Figure 6, the

ROESY spectrum of the ternary system exhibits clear NOE cross-peaks between the H_b protons of the octyl in MVO^{2+} and H_3 and H_5 of α -CD (peaks A and B), which demonstrate that the octyl chain is completely included in the cavity of α -CD. The cross-peak between H_c and H_5 (peak C) further confirms that the primary side of α -CD locates at the portal of MVO^{2+} . Meanwhile, clear NOE cross-peaks between H_a and H_3 (peak D) as well as between the H_x protons of CB[7] and H_2 and H_3 of α -CD¹⁴ provide strong evidence for not only α -CD penetrating into the octyl chain from its secondary side, but also the two macrocycles being closely located in the ternary complex, which suggests hydrogen-bonding interactions between CB[7] and α -CD. That is, CB[7] is moved to the nucleus, and α -CD correspondingly perches on the octyl moiety in $[\text{MVO}\cdot\text{CB}[7]\cdot\alpha\text{-CD}]^{2+}$. Furthermore, we also carried out molecular dynamics simulation¹⁷ on the system, and four snapshots are shown in Figure 7, which are consistent with our experimental results.

Conclusion

In summary, we have presented the controlled wheel-molecular movement on a molecular axle driven by another macroring molecule. The binding constants and the thermodynamic parameters obtained in this study reveal that the exothermic enthalpy changes of the complexation between the axle molecules and α -CD afford an impetus to push CB[7] from the octyl moiety to the bipyridinium moiety of the viologen. This work should not only be useful for supramolecular control of molecular behavior, but also provide insight into designing locomotive molecular devices that are driven by another molecule.

(16) Ong, W.; Kaifer, A. E. *Org. Lett.* **2002**, *4*, 1791.

(17) A 100 ps MD simulation is performed by using the NVT ensemble with a time step of 1 fs at 298 K using the Dreiding force field (Mayo, S. L.; Olafson, B. D.; Goddard, W. A., III. *J. Phys. Chem.* **1990**, *94*, 8897). The cutoff distance is 9 Å with a buffer length of 0.5 Å.




FIGURE 8. Calorimetric titration of CB[7] with MVO²⁺ in Tris buffer (pH 7.4) at 298 K: (a) raw data for sequential 10 μL injections of CB[7] solution into MVO²⁺ solution; (b) “net” heat effect obtained by subtracting the heat of dilution from the heat of reaction, which was analyzed by computer simulation using the one set of binding sites model.

Experimental Section

Materials. All chemicals were of the highest purity commercially available and were used without further purification. BV²⁺ and BnV²⁺ were synthesized according to the refs 18 and 19, respectively. The syntheses of MVO²⁺ and OV²⁺ are described in the Supporting Information.

Isothermal Microcalorimetric Titration. All calorimetric experiments were performed using a thermostated and fully computer

(18) Moon, K.; Kaifer, A. E. *Org. Lett.* **2004**, *6*, 185.

(19) Joseph, J.; Eldho, N. V.; Ramaiah, D. *Chem.—Eur. J.* **2003**, *9*, 5926.

operated VP-ITC calorimeter purchased from a commercial supplier. In this study, all microcalorimetric titrations were performed in an aqueous Tris buffer solution (pH 7.4) at atmospheric pressure and 298 K. Each solution was degassed and thermostated using a ThermoVac accessory before titration. Each microcalorimetric titration experiment consisted of 25 or 29 successive injections. A constant volume (10 μL /injection for α -CD or CB[7]) of host solution (2.87–9.93 mM for each host) in a 0.250 mL syringe was injected into the reaction cell (1.4227 mL) charged with MVO²⁺, OV²⁺, [MVO·CB[7]]²⁺, or [OV·CB[7]]²⁺ solution (0.12–0.32 mM) in the same buffer solution.

A control experiment to determine the heat of dilution was carried out with each run by performing the same number of injections with the same concentration of host compound as used in the titration experiments into a pure buffer solution without the guest compound. The dilution enthalpies determined in control experiments were subtracted from the enthalpies measured in the titration experiments to obtain the net reaction heat.

ORIGIN software, which was used to simultaneously compute the equilibrium constant (K_S) and standard molar enthalpy of reaction (ΔH°) from a single titration curve, gave a standard deviation based on the scatter of the data points in the titration curve. The net reaction heat in each run was calculated by the “one set of binding sites” model unless noted otherwise. Additionally, the first point was removed from the titration curve, acknowledging that the concentration of guest in the cell far exceeded the concentration of the host. A typical titration curve and the fitted results for the complexation of CB[7] with MVO²⁺ are shown in Figure 8. The isothermal calorimetric titration curve obtained by the addition of α -CD to the [OV·CB[7]]²⁺ solution is analyzed by using the equation for the simultaneous formation of the 1:1 and 1:2 complexes (sequential binding sites model).¹⁴ The knowledge of the binding constant (K_S) and molar reaction enthalpy (ΔH°) enabled calculation of the standard free energy of binding (ΔG°) and entropy changes (ΔS°), according to the equation

$$\Delta G^\circ = -RT \ln K_S = \Delta H^\circ - T\Delta S^\circ$$

where R is the gas constant and T is the absolute temperature.

Multiple independent titration runs ($N = 2$) were performed to afford self-consistent thermodynamic parameters, and the averaged values are reported in Table 1. The uncertainties in the thermodynamic parameters reported for host–guest complexation are 2 standard deviations of the mean value.

Acknowledgment. This work was supported by the 973 Program (Grant 2006CB932900), NNSFC (Grant Nos. 90306009, 20421202, and 20673061), and the Special Fund for Doctoral Program from the Ministry of Education of China (Grant 20050055004), which are gratefully acknowledged.

Supporting Information Available: Additional 1D and 2D NMR, ESI-MS, and UV–vis spectra and an ITC titration curve as mentioned in the text. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO0624288